


Complementary and alternative medicine for glycemic control of diabetes mellitus: A systematic review

By Erni Setiyorini

Complementary and alternative medicine for glycemic control of diabetes mellitus: A systematic review

Journal of Public Health Research
2022, Vol. 11(3), 1-10
© The Author(s) 2022
DOI: 10.1177/22799036221106582
journals.sagepub.com/home/phj


Erni Setiyorini¹, Mochammad Bagus Qomaruddin¹, Sony Wibisono²,
Titik Juwariah³, Anggi Setyowati⁴, Ning Arti Wulandari⁵,
Yeni Kartika Sari⁵ and Levi Tina Sari⁵

Abstract

The use of complementary and alternative medicine (CAM) is increasingly popular for the management of diabetes mellitus (DM). The aim of this study was to conduct systematic review of any types of complementary and alternative medicine for glycemic control of diabetes mellitus. Four databases was used in this study, the CINAHL, PUBMED, SCOPUS, and ProQUEST. The systematic review were reported according to the PRISMA guidelines. The keywords were used according to medical subject headings (MeSH) in this study were diabetes mellitus AND complementary and alternative medicine AND blood glucose levels or blood sugar or blood glucose. Articles were limited to 2015-2021 and only in English language. We obtained 231 articles from these databases: CINAHL six articles, PUBMED 85 articles, SCOPUS 66 articles, PROQUEST 74 articles. Then, the final results recorded 17 articles. The results of a systematic review showed the effectiveness of natural products as CAM for glycemic control of DM, namely *Berberis aristata/Silybum marianum*, fenugreek seed, bitter melon, cinnamon or whortleberry supplements, a combination of herbal plants (*C. spinosa*, *R. canina*, and *S. securigera*), *Nigella sativa*, Mulberry juice, chicory, chamomile tea, and bell pepper juice combined with an integrated approach of yoga therapy. Mind body practices such as auditory guided imagery (AGI), qigong and tai chi exercises, and relaxation. Whole system approach, such as acupressure. Health care providers consider CAM for DM management.

Keywords

Complementary and alternative medicine, diabetes mellitus, systematic review

Date received: 19 May 2022; accepted: 19 May 2022

Introduction

Diabetes mellitus (DM) is a chronic disease that happens due to either inadequacy of insulin production by the pancreas or the inability of the body to use insulin effectively. Insulin is hormone that regulate the blood sugar. DM is major cause of heart attack and stroke, and damage the blood vessel of eyes, kidney, and nerves.¹ DM is still worldwide problem.² The prevalence of DM is increased rapidly in developing countries compared to developed countries. By 2019, an estimated 1.5 million deaths were due to this disease and 2.2 million deaths were due to an increase in blood sugar in 2012.³

There are many challenges in the management of diabetes mellitus.³ The first treatment of DM is to use an

antihyperglycemic medicine, but the use of this conventional medicine also has adverse effect.⁴ So that complementary and alternative medicine (CAM) is starting to

¹Public Health Faculty, Universitas Airlangga, Surabaya, Indonesia

²Department of Internal Medicine, Faculty of Medicine, Universitas Airlangga, Surabaya, Indonesia

³Ganesha Husada College of Health, Kediri, Indonesia

⁴School of Nursing, Faculty of Medicine, Universitas Lambung Mangkurat, Banjarbaru, Indonesia

⁵Patria Husada College of Health, Blitar, Indonesia

Corresponding Author:

Erni Setiyorini, Public Health Faculty, Universitas Airlangga, Surabaya, 60115, Indonesia.

Email: emi.setiyorini-2019@fkm.unair.ac.id



become an option for the treatment of chronic diseases such as metabolic disorder.⁵ The National Center for Complementary and Alternative Medicine in the United States stated that CAM is a group of the health system, health practice, and health product considered as not part of conventional medicine. Complementary medicine is used with conventional therapy whereas alternative medicine is used as a substitute for conventional therapy.⁶ More than 400 plants and substances are evaluated for DM treatment especially DM type 2.⁷

The use of CAM has become increasingly popular for the treatment of DM in that it is related to minimal complication and minimal cost. Moreover, CAM is become an option considering that cultural and psychosocial factor,⁸ health belief⁹ as well as value related to religion.¹⁰ The survey in Canada about the use of alternative remedies showed that 44% were taking supplements and 31% were taking alternative medications.¹¹ The percentage of diabetic patients who used CAM in United States is larger than that in Australia with 57%¹² and 25% consecutively.¹³ Furthermore, people in various countries are still dependent on CAM for treatment such as Asia and Africa.¹⁴

This study aims to review the most-common CAM used to control glycemic in patients with diabetes mellitus. CAM intervention for DM can be divided into two major categories namely mind-body practice (including meditation, relaxation, and aromatherapy) and natural products (including herbal medicines, vitamins, minerals, and supplements). Additionally, there are some of whole system alternative medicine approach (including traditional Chinese medicine, acupuncture, and reflexology).¹⁵

Research methods

Literature search strategy

This article used a systematic review design and followed previous study to guide the review,¹⁶ the PRISMA guidelines was used to report the systematic review.¹⁷ The study using CINAHL, PUBMED, SCOPUS, and PROQUEST database from January 2015 to September 2019. The keywords were used according to medical subject headings (MeSH) in this study were "diabetes mellitus AND complementary and alternative medicine AND blood glucose levels or blood sugar or blood glucose." Tables were created to assist the result of this study.

Inclusion criteria

We searched original studies and already published it. Exclusion articles were article not in the English language, not full paper article, not in human, and not master or dissertation.³ The population in this study was individuals with Type 1 diabetes mellitus (T1DM), Type 2 diabetes mellitus (T2DM), and gestational diabetes mellitus,¹⁸ the

intervention was complementary and alternative medicine, the type of study was randomized controlled trials (RCT), and the outcome was glycemic control. Observational studies, non-randomize control, and case control studies were not included in this review.

Intervention

Complementary and alternative medicine was categorized into mind-body practices (e.g. hypnosis, CBT, relaxation, biofeedback, meditation, aromatherapy), natural products (e.g. herbs, vitamins, minerals, and supplements) and whole system approaches (e.g. traditional Chinese medicine, reflexology, acupuncture, homeopathy).¹⁵ The intervention used in the study was one of these interventions or a combination.

Outcome

The original articles included in the study was glycemic control or diabetes mellitus patients, such as lowering A1C blood glucose level, fasting blood glucose, random blood glucose, or oral glucose tolerance testing.¹⁹

Risk of bias

Two independent authors screened 21 articles. The research methodology was evaluated using the JBI Critical Appraisal checklist.²⁰ The checklist consisted of questions that were used to appraise the original articles. Scores less than 50% were excluded from the study to avoid bias (Table 1).

Results

Figure 1 showed the data extraction process. We obtained 231 articles from these databases: CINAHL six articles, PUBMED 85 articles, SCOPUS 66 articles, PROQUEST 74 articles. Then, we removed duplicate article six articles. We screened on title and abstract, then we got 37 eligible articles. Further, we exclude the articles that did not include in our criteria, such as population, intervention, study type, and outcome. Seventeen articles were included in our systematic review.

Natural products (herbal products, vitamins, and supplements)

Natural products in the systematic review were *Berberis aristata/Silybum marianum*, fenugreek seed, bitter melon, cinnamon or whortleberry supplements, a combination of herbal plants; *C. spinosa*, *R. canina*, and *S. securigera*, *Nigella sativa*, mulberry juice, chicory, chamomile tea, and also bell pepper juice combined with the integrated approach of yoga therapy (IAYT) (Table 2).

Table 1. Risk of bias.

Title	Criteria (checklist "√")													Value, %
	1	2	3	4	5	6	7	8	9	10	11	12	13	
7 Derosa et al. ²¹	√	√	√	√	√	√	√	√	√	√	√	√	√	100
Hadi et al. ²²	√	√	√	√	√	√	√	√	√	√	√	√	√	100
Kim et al. ²³	√	√	√	√	√	-	√	√	√	√	√	√	√	92
Mehrzadi et al. ²⁴	√	√	√	√	√	√	√	√	√	√	√	√	√	100
Mirfeizi et al. ²⁵	√	√	√	√	√	√	√	√	√	√	√	√	√	100
Moraes et al. ²⁶	√	√	√	√	√	√	√	√	√	√	√	√	√	100
Moustafa et al. ²⁷	√	-	√	-	-	-	√	√	√	√	√	√	√	69
Nagasukeerthi et al. ³⁰	√	√	√	√	√	-	√	√	√	√	√	√	√	92
Riche et al. ²⁸	√	√	√	√	√	-	√	√	√	√	√	√	√	92
Zemestani et al. ²⁹	√	√	√	√	-	-	√	√	√	√	√	√	√	84
El-Shamy et al. ³⁴	√	√	√	√	√	√	√	√	√	√	√	√	√	100
17 Kullah ³⁵	√	√	√	√	-	-	√	√	√	√	√	√	√	84
Kumar et al. ³⁶	√	√	√	√	√	√	√	√	√	√	√	√	√	100
Mooventhan et al. ³⁷	√	√	√	√	√	√	√	√	√	√	√	√	√	100
Gelernter et al. ³¹	√	√	√	√	√	√	√	√	√	√	√	√	√	100
Li et al. ³²	√	√	√	√	-	√	√	√	√	√	√	√	√	92
Paschali et al. ³³	√	-	√	-	-	-	√	√	√	√	√	√	√	69

1. Was true randomization used for assignment of participants to treatment groups?, 2. Was allocation to treatment groups concealed?, 3. Were treatment groups similar at the baseline?, 4. Were participants blind to treatment assignment?, 5. Were those delivering treatment blind to treatment assignment?, 6. Were outcomes assessors blind to treatment assignment?, 7. Were treatment groups treated identically other than the intervention of interest?, 8. Was follow up complete and if not, were differences between groups in terms of their follow up adequately described and analyzed?, 9. Were participants analyzed in the groups to which they were randomized?, 10. Were outcomes measured in the same way for treatment groups?, 11. Were outcomes measured in a reliable way?, 12. Was appropriate statistical analysis used?, 13. Was the trial design appropriate, and any deviations from the standard RCT design (individual randomization, parallel groups) accounted for in the conduct and analysis of the trial?

The results of the study showed that *25* *erberis aristata*/*Silybum marianum* had positive effect on glycaemic control in type 1 diabetes mellitus (T1DM) compared to the control group, one tablet at lunch and one tablet at dinner, for 6 months.²¹ *Fenugreek seed* (FS) has beneficial to reduce fasting blood sugar *22* in type 2 DM (T2DM) compared to the control group. 5 g FS powder was consumed mixed with water, three times daily, for 8 weeks.²² *29* Bitter melon has positive effect to reduce blood sugar in patients with type 2 DM (T2DM) compared to the control group who *1* received a placebo capsule. Bitter melon was consumed twice a day for 12 weeks.²³ The combination of herbal plants, *C. spinosa*, *R. canina* and *S. securigera* was as effective as glycaemic control in type 2 DM (T2DM) patients. This combination was consumed two tablets, once a day.²⁴

The use of cinnamon or whortleberry *3* supplements as an additional treatment could reduce blood sugar levels in type 2 DM patients. This study divided respondents into three groups, the cinnamon group, whortleberry group, and placebo groups. Respondents received cinnamon (1000 mg/day) or Caucasian whortleberry (1000 mg/day) for 3 months. Blood sugar levels decreased in the cinnamon or whortleberry supplement group compared to the placebo group.²⁵ Chicory was also used as a supplement for type 2 DM patients and had effect to decrease fasting blood sugar. Chicory was consumed 10 g daily for 2 months.²⁶

1 Nigella sativa oil 1350 mg/day, for 3 months in newly diagnosed with type 2 DM patients had effect to decrease fasting blood sugar, 2 h post-prandial, and A1C compared *30* metformin.²⁷ Mulberry leaves (1000 mg) was given three times a day with meals, for 3 months had effect to decrease post prandial blood sugar levels compared to placebo in type 2 DM patients.²⁸ Chamomile tea has an effect as a glycaemic control (3 g/1 *18* ml with hot water) among type 2 DM patients, three times a day after meals for 8 weeks compared to a control group.²⁹

Bell pepper juice with the integrated approach of yoga therapy (IAYT) for four consecutive days *1* had effect to reduce post prandial blood sugar levels compared to the control group who only received the integrated approach of yoga therapy (IAYT) in type 2 DM patients.³⁰

Mind body practices

Mind body practices in this study were auditory guided imagery (AGI), qigong and tai chi exercises, and relaxation. The study showed that auditory guided imagery (AGI) accompanied by music for 5 days with a duration of 7 min, two times a day, had effect to reduce blood sugar in children with type 1 DM (T1DM).³¹ Another study was qigong and tai chi exercise for 12 weeks. This study divided groups into three groups; qigong, tai chi exercise, and placebo. The results of the study showed




Figure 1. Flowchart of the study selection.

that qigong had better effect on reducing fasting blood sugar in type 2 DM (T2DM) patients.³² Another study showed that relaxation had effect to reduce fasting blood sugar in type 1 DM (T1DM) patients compared to the placebo group. The intervention was carried out for 8 weeks (Table 3).³³

Whole system approach

The whole system approach in this study is acupressure at certain points. The study showed that acupressure on ST36, BL23, and BL13 for 12 weeks, 3 min, three sessions/week had effect to reduce blood sugar levels in gestational diabetes mellitus compared to the control group who only performed antenatal care.³⁴ Another study showed that acupressure on Zusanli point (ST-36) for 30 min had effect to reduce blood sugar in diabetes mellitus.³⁵

Another study also showed that acupressure on CV-12 (Zhongwan) for 30 min. This study showed that 30 min of CV-12 had effect to reduce blood sugar among type 2 DM patients compared to the placebo group that received stimulated at the right side of the abdomen (1-cun in

beside the CV-12).³⁶ Furthermore, acupressure on Zusanli (Zusanli) for 30 min was also effective in lowering blood sugar compared to the control group at the placebo point (Table 4).³⁷

Discussion

The aim of this study was to conduct a systematic review the most-common CAM used to control glycemic in patients with diabetes mellitus published in the last 5 years ranging from 2015 to 2021. Seventeen articles are included to investigate the effect of CAM to control blood sugar in patients with diabetes mellitus, including natural products, mind-body practice, and whole system approach.

Natural product

Natural products, such as medicinal plants and supplements, are the product widely used in CAM for patients with DM. This is aligned with a previous study which states that natural product is the popular complementary health in the United States.³⁸

Table 2. Systematic reviews of natural product for diabetes mellitus published since January 2015 until September 2021.

Author	Country	Intervention evaluated	Condition treated	Number of studies	Study design	Conclusion	Mention of adverse effects
Derosa et al. ²¹	Italia	<i>Berberis aristata/Silybum marianum</i> 588/105 mg	Diabetes mellitus Type 1	85	RCT	There was a decrease of FPG, and PPG with <i>B. aristata/S. marianum</i> both compared to baseline and placebo	Yes
Hadi et al. ²²	Iran	Fenugreek seed (FS)	Diabetes mellitus Type 2	50	RCT	FS consumption resulted in a significant decrease in fasting plasma glucose (20 G)	Yes
Kim et al. ²³	Korea	<i>Momordica charantia</i> (bitter melon)	Diabetes mellitus Type 2	90	RCT	the average fasting glucose level of the bitter melon group decreased	Yes
Mehrzadi et al. ²⁴	Iran	6 additional herbal <i>Capparis spinosa, Rosacannina, Securidaca securigera, Silybum marianum, Urtica dioica, Trigonella foenum-graecum</i> and <i>Vaccinium arctostaphylos</i>	Diabetes mellitus Type 2	150	RCT	the fasting plasma glucose, HbA1c in herbal combination were decreased significantly	Yes
Mirfeizi et al. ²⁵	Iran	cinnamon or whortleberry supplements	Diabetes mellitus Type 2	105	RCT	the use of cinnamon and whortleberry in addition to conventional medical treatment is recommended to adjust weight and blood glucose levels in patients with T2DM	Yes
Moraes et al. ²⁶	Iran	chicory inulin supplement	Type 2 diabetic mellitus (T2DM)	46	RCT	Significant reductions in fasting serum glucose (FSG), Hb A1C, AST and ALP concentrations were observed in chicory-treated group.	No
Moustafa et al. ²⁷	Egypt	<i>Nigella sativa</i>	Type 2 diabetic mellitus (T2DM)	66	RCT	NS oil administration at a dose of 1350 mg per day in newly diagnosed patients with type 2 diabetes mellitus was inferior to metformin in terms of lowering FBG, 2 h pp, A 38 %B	Yes
Nagasukeerthi et al. ³⁰	India	13 Bell pepper (<i>Capsicum annum var. grossum</i>) juice with integrated approach of yoga therapy	Type 2 diabetic mellitus (T2DM)	50	RCT	a significant reduction in Post prandial blood glucose (PPBG), was observed in the study group	Yes
Riche et al. ²⁸	USA	Mulberry leaves	Type 2 diabetic mellitus (T2DM)	24	RCT	Post-prandial SMBG was significantly decreased at 3 months in the MLE group	Yes
Zemestani et al. ²⁹	Iran	Chamomile tea	Type 2 diabetes mellitus (T2DM)	64	RCT	9 rsus baseline that short term intake of chamomile tea had beneficial effects on glycemic control and antioxidant status	Yes

Table 3. Systematic reviews of mind-body practices for diabetes mellitus published since January 2015 until September 2021.

Author	Country	Intervention evaluated	Condition treated	Number of studies	Study design	Conclusion	Mention of adverse effects
Gelernter et al. ³¹	Israel	Auditory guided imagery (AGI) accompanied by background music and background music solely (BMS)	Diabetes mellitus Type 1	13 children	RCT	Adding AGI sessions of 7 min, to the multidisciplinary management of pediatric population with T1DM may contribute to a decrease in short-term glucose concentration	No
Li et al. ³²	China	Qigong and tai chi exercises	Diabetes mellitus Type 2	103	RCT	there was a significant negative correlation between the duration of T2DM and the relative changes in FPG levels after qigong intervention	Yes
Paschali et al. ³³	Greece	Relaxation training	Diabetes mellitus Type 1 (T1DM)	46	RCT	the main metabolic measurement of blood glucose levels and HbA1C revealed significant differences over time	No

Berberis aristata/Silybum marianum. *Berberis aristata/Silybum marianum* is medicinal herbs originated from Asia and Southern Europe, but now it can be found in many countries around the world.³⁹ This plant contains antioxidant. In this literature review, *Berberis aristata* can be added with insulin therapy as a glycemic control for patients with DM type.⁴⁰

Fenugreek (*Trigonella foenum-graecum*). Fenugreek is an herb that has a special scent and grows in India and some countries in South Africa.⁴¹ Fenugreek contains alkaloids and steroidal saponins which can be used as an antidiabetic agent.⁴² This literature review uses fenugreek seeds to decrease fasting blood sugar.²²

³⁹ Bitter melon (*Momordica charantia*)

Bitter melon has been frequently used as a therapy for decreasing blood sugar in diabetes mellitus.²⁷ Bitter melon, a member of the Curcubitaceae family, is a plant that grows in tropical areas such as Amazon Basin, Africa, Asia, the Caribbean, and Southern America. It has green leaves, yellow flowers, and its fruit like a cucumber. Bitter melon extract has a similar structure to animal insulin. The frequent adverse effects are hypoglycemia and dizziness,⁴³ while in this study there is no serious side effects when the intervention was conducted.²³

Whortleberry (*Vaccinium arctostaphylos* L.). The Iranian people widely used whortleberry as a traditional medicine.⁴⁴

It has been reported to have antioxidant, anti-cancer, anti-inflammatory, and anti-hyperglycemia properties.^{45,46}

Cinnamon (*Cinnamomum*). Cinnamon is a sweet spice that is commonly used in Greece and Rome. This plant is stated in Bible and Chinese texts. Cinnamon, originated from Sri Lanka and part of India, is a tropical plant that has affects blood sugar, body mass index (BMI)⁴⁷ and insulin.⁴⁸

Chicory inulin. Chicory is an agricultural crop from continental Europe. This herb has been widely consumed in various countries in Western and Eastern countries.⁴⁹ Chicory has been reported to have function as a prebiotic,⁵⁰ and to decrease blood sugar level.⁵¹ This article review found that chicory inulin could lower the level of fasting blood glucose.²⁶

Nigella sativa. *Nigella sativa*, an original plant from Mediterranean to Western Asia and Northern India, is widely produced in India, Bangladesh, Nepal, Sri Lanka, Iraq, and Pakistan.⁴⁸ This plant contains active substances known as thymoquinone and has antioxidant effect.⁵² In addition, *Nigella sativa* can reduce appetite and weight, and glycemic control.⁵³

Bell pepper (*Capsicum annuum* L.). Bell pepper (*Capsicum annuum* L., also known as chili pepper, has been used by native Americans. The essential substance of this plant is an alkaloid known as capsinoids. It has been reported that bell pepper is the main source of vitamin C. Also, bell pepper contains antioxidant.⁵⁴ In this research, bell pepper

Table 4. Systematic reviews of whole system approaches for diabetes mellitus published since January 2015 until September 2021.

Author	Country	Intervention evaluated	Condition treated	Number of studies	Study design	Conclusion	Mention of adverse effects
El-Shamy et al. ³⁴	Egypt	Acupressure	Gestational diabetes mellitus	30 female	RCT	5 After 12 weeks intervention had shown that 75 g oral glucose tolerance test (OGTT), insulin resistance, number of required insulin and measure of utilized insulin were significantly reduced	No
Fitrullah and Rousdy ³⁵	Indonesia	Acupressure	Diabetes mellitus	30	RCT	5 Acupressure at the Zusanli (ST 36) acu-point can lower blood glucose levels significantly	Yes
Kumar et al. ³⁶	India	Acupuncture	Diabetes mellitus Type 2	40	RCT	A 50% significant reduction in random blood glucose level in Acupuncture group compared to its baseline	No
Mooventhan et al. ³⁷	India	Acupuncture ST 36	Diabetes mellitus Type 2	60	RCT	14 present study showed a significant reduction in random blood glucose levels in the acupuncture group compared to the placebo control group	No

was combined with yoga so that it can be used as a management of DM type 2.³⁰

Mulberry leaves. Mulberry is commonly grown in the mountain of the tropical area for instance Asia, Middle East, and Southern America.⁵⁵ It leaves has an advantage for the treatment of DM type 2. Mulberry leaves can be a useful complementary food to lower postprandial blood sugar according to Riche et al.²⁸

Chamomile tea. Chamomile, having the Latin name *Chamomilla recutita L.*, is a medicinal herb from Europe and Western Asia. Chamomile flowers are commonly used as tea for medicinal purpose.⁵⁶ The result of the research by Zemestani et al.²⁹ found that chamomile has anti-hyperglycemia and antioxidant properties.

Traditional herb.

6 Several herbal medicine combinations consisting of *Capparis spinosa*, *Rosa canina*, *Securidaca securigera*, *Silybum marianum*, *Urtica dioica*, *Trigone* 1 *foenum-graecum*, and *Vaccinium arctosphylos* is used to glycemic control in patients with DM type 2, and there was not found any adverse effect.²⁴

Mind-body practice

Mind-body practice focus on physic and psychological approach for instance relaxation (breathing exercise and

guided imagery), tai chi, yoga, and music therapy. Auditory guided imagery (AGI), qigong and tai chi, and relaxation are used in this review.

Auditory guided imagery (AGI). Auditory guided ima 4 ty is a psychological intervention by listening to relaxing music and verbally described on images formed in the mind in order to feel relaxed and focus.⁵⁷ Moreover, the feeling of relaxation that appeared 36 auditory guided imagery has a glycemic control effect and improves the quality of life in patients with DM type 12 the result of this review stated that AGI has a decrease of blood glucose effect in children with DM type 2 at a short period of time.³¹

Qigong and Tai Chi. Psychical activity is needed to manage blood glucose in diabetic patient.⁵⁸ Qigong is a breathing exercise and is an essential part of Chinese medicine.⁵⁹ This exercise affects blood sugar.⁶⁰ Tai chi 33 originated from China and slow-motion such as dance and integrates musculoskeletal, breathing, and meditation.⁶¹ Tai chi can be used as a chronic DM management since this exercise can alleviate fasting blood glucose.⁶² This review found that qigong is more effective for managing patients with DM type 2 than tai chi.³²

Relaxation. S 45 triggers chronic hyperglycemia. Relaxation is known to reduce blood glucose levels so that it can be used for treating patients with DM type.³³

Whole system approach

The whole system approach in this study is acupressure at several points. Acupressure has been practiced in Traditional Chinese Medicine (TCM) for 5000 years. Acupressure involves pressing several points on the body. This technique can stimulate the production of endorphins in the brain, relieve pain, and enhance comfort.⁶³ The result of the review showed that acupressure could be used as glycemic control in patients with gestational diabetes mellitus³⁴ and diabetes mellitus type 2.^{35–37}

Limitation of the study

The articles of the systematic review still have some limitations, such as the sample in the articles was still too small and duration of the study and follow up. However, the articles in this study mentioned the adverse effects. Further study is needed to conduct study related to the potential interaction of CAM therapy with medicine

Conclusion

CAM for glycemic control of diabetes mellitus are natural product, mind body practice, and whole system approach. The results of the systematic review showed that natural products were the most widely used type for glycemic control of DM, but the side effect must be considered when using these natural products.

Acknowledgements

We would like to thank LPDP for their support of this study.

Author contributions

The authors contributed equally.

Declaration of conflicting interests

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Funding

The author(s) disclosed receipt of the following financial support for the research, authorship, and/or publication of this article: The work was supported by LPDP Indonesia.

Ethics approval

This study is systematic review and the protocol of this study was registered at Prospero number CRD42021287919.

Significance for public health

Complementary and alternative medicines has been increasingly and become public interest. CAM practice have emphasized health promotion and being a part of public health. In addition CAM practice is also increasingly popular for the management of diabetes mellitus (DM) due to minimal complication and minimal

cost. Those, the aim of this study was to conduct systematic review of any types of complementary and alternative medicine for glycemic control of diabetes mellitus. The result of this study provided the evidence for health care provider for using CAM for glycemic control.

References

1. WHO. Diabetes 2021, <https://www.who.int/news-room/fact-sheets/detail/diabetes> (accessed 29 June 2022).
2. Reddy SS. Health outcomes in type 2 diabetes. *Int J Clin Pract Suppl* 2000; 2000(113): 46–53.
3. Xie W, Zhao Y and Zhang Y. Traditional Chinese medicines in treatment of patients with type 2 diabetes mellitus. *Evid Based Complement Alternat Med* 2011; 2011: 726723.
4. Philippe J and Raccach D. Treating type 2 diabetes: how safe are current therapeutic agents? *Int J Clin Pract* 2009; 63(2): 321–332.
5. Bausell RB, Lee W-L and Berman BM. Demographic and health-related correlates to visits to complementary and alternative medical providers. *Med Care* 2001; 39: 190–196.
6. Kumar D, Bajaj S and Mehrotra R. Knowledge, attitude and practice of complementary and alternative medicines for diabetes. *Public Health* 2006; 120(8): 705–711.
7. Chang CL, Lin Y, Bartolome AP, et al. Herbal therapies for type 2 diabetes mellitus: chemistry, biology, and potential application of selected plants and compounds. *Evid Based Complement Alternat Med* 2013; 2013: 378657.
8. Astin JA. Why patients use alternative medicine: results of a national study. *JAMA* 1998; 279(19): 1548–1553.
9. Islahudin F, Shahdan IA and Mohamad-Samuri S. Association between belief and attitude toward preference of complementary alternative medicine use. *Patient Prefer Adherence* 2017; 11: 913–918.
10. Lee GB, Cham TC, Chew ZH, et al. Complementary and alternative medicine use in patients with chronic diseases in primary care is associated with perceived quality of care and cultural beliefs. *Fam Pract* 2004; 21(6): 654–660.
11. Ryan EA, Pick ME and Marceau C. Use of alternative medicines in diabetes mellitus. *Diabetic Med* 2001; 18(3): 242–245.
12. Yeh GY, Eisenberg DM, Davis RB, et al. Use of complementary and alternative medicine among persons with diabetes mellitus: results of a national survey. *Am J Public Health* 2002; 92(10): 1648–1652.
13. Tan AC and Mak JC. Complementary and alternative medicine in diabetes (CALMIND): a prospective study. *J Complement Integr Med* 2015; 12(1): 95–99.
14. Peltzer K and Pengpid S. Prevalence and determinants of traditional, complementary and alternative medicine provider use among adults from 32 countries. *Chin J Integr Med* 2018; 24(8): 584–590.
15. NCCIH. Complementary, alternative, or integrative health: what's in a name?, <https://www.nccih.nih.gov/health/complementary-alternative-or-integrative-health-whats-in-a-name> (2021, accessed 29 September 2021).
16. Grossman LD, Roscoe R and Shack AR. Complementary and alternative medicine for diabetes. *Can J Diabetes* 2018; 42(Suppl 1): S154–S161.

17. PRISMA. Welcome to the preferred reporting items for systematic reviews and meta-analyses (PRISMA) website!, <https://prisma-statement.org/> (accessed 29 June 2022).
18. World Health Organization. *Classification of diabetes mellitus*. Geneva: World Health Organization, 2019.
19. Patel P and Macerollo A. Diabetes mellitus: diagnosis and screening. *Am Fam Phys* 2010; 81(7): 863–870.
20. Tufanaru CMZ, Aromataris E, Campbell J, et al. Systematic reviews of effectiveness. In: Aromataris E and Munn Z (eds) *JBI manual for evidence synthesis*. <https://synthesismanual.jbi.global/> (2020, accessed 29 June 2022).
21. Derosa G, D'Angelo A and Maffioli P. The role of a fixed Berberis aristata/Silybum marianum combination in the treatment of type 1 diabetes mellitus. *Clin Nutr* 2016; 35(5): 1091–1095.
22. Hadi A, Arab A, Hajianfar H, et al. The effect of fenugreek seed supplementation on serum irisin levels, blood pressure, and liver and kidney function in patients with type 2 diabetes mellitus: a parallel randomized clinical trial. *Complement Ther Med* 2020; 49: 102315.
23. Kim SK, Jung J, Jung JH, et al. Hypoglycemic efficacy and safety of *Momordica charantia* (bitter melon) in patients with type 2 diabetes mellitus. *Complement Ther Med* 2020; 52: 102524.
24. Mehrzadi S, Mirzaei R, Heydari M, et al. Efficacy and safety of a traditional herbal combination in patients with type II diabetes mellitus: a randomized controlled trial. *J Diet Suppl* 2021; 18(1): 31–43.
25. Mirfeizi M, Mehdizadeh Tourzani Z, Mirfeizi SZ, et al. Controlling type 2 diabetes mellitus with herbal medicines: a triple-blind randomized clinical trial of efficacy and safety. *J Diabetes* 2016; 8(5): 647–656.
26. Moraes MR, Simões HG, Farhangi MA, et al. The effect of enriched chicory inulin on liver enzymes, calcium homeostasis and hematological parameters in patients with type 2 diabetes mellitus: a randomized placebo-controlled trial. *PLoS One* 2016; 10(4): 265–271.
27. Moustafa HAM, El Wakeel LM, Halawa MR, et al. Effect of *Nigella sativa* oil versus metformin on glycemic control and biochemical parameters of newly diagnosed type 2 diabetes mellitus patients. *Endocrine* 2019; 65(2): 286–294.
28. Riche DM, Riche KD, East HE, et al. Impact of mulberry leaf extract on type 2 diabetes (Mul-DM): a randomized, placebo-controlled pilot study. *Complement Ther Med* 2017; 32: 105–108.
29. Zemestani M, Rafrat M and Asghari-Jafarabadi M. Chamomile tea improves glycemic indices and antioxidant status in patients with type 2 diabetes mellitus. *Nutrition* 2016; 32(1): 66–72.
30. Nagasukeerthi P, Mooventhan A and Manjunath NK. Short-term effect of add on bell pepper (*Capsicum annum var. grossum*) juice with integrated approach of yoga therapy on blood glucose levels and cardiovascular functions in patients with type 2 diabetes mellitus: a randomized controlled study. *Complement Ther Med* 2017; 34: 42–45.
31. Gelernter R, Lavi G, Yanai L, et al. Effect of auditory guided imagery on glucose levels and on glycemic control in children with type 1 diabetes mellitus. *J Pediatr Endocrinol Metab* 2016; 29(2): 139–144.
32. Li X, Si H, Chen Y, et al. Effects of fitness qigong and tai chi on middle-aged and elderly patients with type 2 diabetes mellitus. *PLoS One* 2020; 15(12): e0243989.
33. Paschali AA, Peppou LE and Benroubi M. Relaxation training significantly reduced blood glucose levels in patients with type 1 diabetes mellitus. *Hormones* 2020; 19(2): 215–222.
34. El-Shamy FF, El-Kholy SS, Labib M, et al. Ameliorative potential of acupressure on gestational diabetes mellitus: a randomized controlled trial. *J Complement Integr Med* 2019; 16(1): 20180011.
35. Fitrullah and Rousdy A. Effectiveness of acupressure at the Zusanli (ST-36) acupoint as a comfortable treatment for diabetes mellitus: a pilot study in Indonesia. *J Acupunct Meridian Stud* 2017; 10(2): 96–103.
36. Kumar R, Mooventhan A and Manjunath NK. Immediate effect of needling at CV-12 (Zhongwan) acupuncture point on blood glucose level in patients with type 2 diabetes mellitus: a pilot randomized placebo-controlled trial. *J Acupunct Meridian Stud* 2017; 10(4): 240–244.
37. Mooventhan A, Ningombam R and Nivethitha L. Effect of bilateral needling at an acupuncture point, ST-36 (Zusanli) on blood glucose levels in type 2 diabetes mellitus patients: a pilot randomized placebo controlled trial. *J Complement Integr Med* 2020; 17(3): 20190100.
38. Su D and Li L. Trends in the use of complementary and alternative medicine in the United States: 2002–2007. *J Health Care Poor Underserved* 2011; 22(1): 296–310.
39. Bahmani M, Shirzad H, Rafieian S, et al. *Silybum marianum*: beyond hepatoprotection. *J Evid Base Compl Altern Med* 2015; 20(4): 292–301.
40. Zhou J, Zhou S, Tang J, et al. Protective effect of berberine on beta cells in streptozotocin- and high-carbohydrate/high-fat diet-induced diabetic rats. *Eur J Pharmacol* 2009; 606(1–3): 262–268.
41. Srinivasan K. Fenugreek (*Trigonella foenum-graecum*): a review of health beneficial physiological effects. *Food Rev Int* 2006; 22(2): 203–224.
42. Gong J, Fang K, Dong H, et al. Effect of fenugreek on hyperglycaemia and hyperlipidemia in diabetes and prediabetes: a meta-analysis. *J Ethnopharmacol* 2016; 194: 260–268.
43. Basch E, Gabardi S and Ulbricht C. Bitter melon (*Momordica charantia*): a review of efficacy and safety. *Am J Health Syst Pharm* 2003; 60(4): 356–359.
44. Fathi M, Naghdi Badi H, Ghanbari A, et al. Pomological and phytochemical diversity in Iranian populations of Caucasian whortleberry (*Vaccinium arctostaphylos L.*). *Sci Hortic* 2019; 243: 107–115.
45. Hafizur RM, Kabir N and Chishti S. Modulation of pancreatic β -cells in neonatally streptozotocin-induced type 2 diabetic rats by the ethanolic extract of *Momordica charantia* fruit pulp. *Nat Prod Res* 2011; 25(4): 353–367.
46. Takikawa M, Inoue S, Horio F, et al. Dietary anthocyanin-rich bilberry extract ameliorates hyperglycemia and insulin sensitivity via activation of AMP-activated protein kinase in diabetic mice. *Nutr J* 2010; 140(3): 527–533.
47. Nasri H, Madihi Y and Marikhi A. Commentary on: effects of cinnamon consumption on glycemic status, lipid profile and body composition in type 2 diabetic patients. *Int J Prev Med* 2013; 4(5): 618–619.

48. Caquet R. Antioxidant properties of spices, herbs and other sources. New York Springer science, Business Media: tailored for a challenging environment. *Microbiology and molecular biology reviews* 2004; 70(2): 344.
49. Lucchin M, Varotto S, Barcaccia G, et al. Chicory and endive. In: Prohen J and Nuez F (eds) *Vegetables I*. New York, NY: Springer, 2008. pp.3–48.
50. de Wiele TV, Boon N, Possemiers S, et al. Prebiotic effects of chicory inulin in the simulator of the human intestinal microbial ecosystem. *FEMS Microbiol Ecol* 2004; 51(1): 143–153.
51. Ning C, Wang X, Gao S, et al. Chicory inulin ameliorates type 2 diabetes mellitus and suppresses JNK and MAPK pathways in vivo and in vitro. *Mol Nutr Food Res* 2017; 61(8): 1600673.
52. Javanbakht J, Hobbenaghi R, Hosseini E, et al. Histopathological investigation of neuroprotective effects of *Nigella sativa* on motor neurons anterior horn spinal cord after sciatic nerve crush in rats. *Pathol Biol* 2013; 61(6): 250–253.
53. Alimohammadi S, Hobbenaghi R, Javanbakht J, et al. Protective and antidiabetic effects of extract from *Nigella sativa* on blood glucose concentrations against streptozotocin (STZ)-induced diabetic in rats: an experimental study with histopathological evaluation. *Diagn Pathol* 2013; 8(1): 137–7.
54. Nadeem M, Anjum FM, Khan MR, et al. Antioxidant potential of bell pepper (*Capsicum annum L.*): a review. *Pak J Food Sci* 2011; 21(1–4): 45–51.
55. Ohyama K and Oka S. *Mulberry: cell and tissue culture in forestry*. Dordrecht: Springer, 1987. pp.272–284.
56. McKay DL and Blumberg JB. A review of the bioactivity and potential health benefits of chamomile tea (*Matricaria recutita L.*). *Phytother Res: Int J Pharmacol Toxicol Eval Nat Prod Der* 2006; 20(7): 519–530.
57. Rossi EL. *The psychobiology of mind-body healing: new concepts of therapeutic hypnosis*. New York, NY: WW Norton & Company, 1993.
58. Zinman B, Ruderman N, Campaigne BN, et al. Physical activity/exercise and diabetes mellitus. *Diabetes Care* 2003; 26(Suppl 1): S73–S77.
59. Liu T. *Chinese medical qigong*. London: Singing Dragon, 2010.
60. Lee MS, Chen KW, Choi T-Y, et al. Qigong for type 2 diabetes care: a systematic review. *Complement Ther Med* 2009; 17(4): 236–242.
61. Price CJ and Thompson EA. Measuring dimensions of body connection: body awareness and bodily dissociation. *J Altern Complement Med* 2007; 13(9): 945–953.
62. Lee MS, Jun JH, Lim HJ, et al. A systematic review and meta-analysis of tai chi for treating type 2 diabetes. *Maturitas* 2015; 80(1): 14–23.
63. Yang M-H and Lin L-C. [Acupressure in the care of the elderly]. *Hu li za zhi J Nurs* 2007; 54(4): 10–15.

Complementary and alternative medicine for glycemic control of diabetes mellitus: A systematic review

ORIGINALITY REPORT

19%

SIMILARITY INDEX

PRIMARY SOURCES

1	link.springer.com Internet	82 words — 2%
2	repository.unair.ac.id Internet	74 words — 2%
3	www.ncbi.nlm.nih.gov Internet	52 words — 1%
4	www.degruyter.com Internet	47 words — 1%
5	cybermedlife.eu Internet	43 words — 1%
6	pubmed.ncbi.nlm.nih.gov Internet	43 words — 1%
7	watermark.silverchair.com Internet	38 words — 1%
8	journal.ipm2kpe.or.id Internet	18 words — < 1%
9	Maryam Zemestani, Maryam Rafraf, Mohammad Asghari-Jafarabadi. "Chamomile tea improves	16 words — < 1%

glycemic indices and antioxidants status in patients with type 2 diabetes mellitus", Nutrition, 2016

Crossref

10 16 words — < 1%

bmccomplementmedtherapies.biomedcentral.com

Internet

11 16 words — < 1%

www.mdpi.com

Internet

12 13 words — < 1%

"Textbook of Diabetes", Wiley, 2010

Crossref

13 13 words — < 1%

rsdjournal.org

Internet

14 12 words — < 1%

Muhammed Shabil, Ganesh Bushi, Pavan Kalyan Bodige, Pavan Sagar Maradi et al. "Effect of Fenugreek on Hyperglycemia: A Systematic Review and Meta-Analysis", Medicina, 2023

Crossref

15 12 words — < 1%

bugsigdb.org

Internet

16 12 words — < 1%

www.pubfacts.com

Internet

17 11 words — < 1%

"Supportive Cancer Care with Chinese Medicine", Springer Science and Business Media LLC, 2010

Crossref

18 11 words — < 1%

M. Rafraf, M. Zemestani, M. Asghari-Jafarabadi. "Effectiveness of chamomile tea on glycemic control and serum lipid profile in patients with type 2 diabetes", Journal of Endocrinological Investigation, 2014

-
- 19 medcraveonline.com 11 words — < 1%
Internet
-
- 20 www.researchgate.net 11 words — < 1%
Internet
-
- 21 www.sysrevpharm.org 11 words — < 1%
Internet
-
- 22 Amir Hadi, Arman Arab, Hossein Hajianfar, Behrouz Talaei et al. "The effect of fenugreek seed supplementation on serum irisin levels, blood pressure, and liver and kidney function in patients with type 2 diabetes mellitus: A parallel randomized clinical trial", *Complementary Therapies in Medicine*, 2020 10 words — < 1%
Crossref
-
- 23 Anna Surgean Veterini, Nancy Margarita Rehatta, Annis Catur Adi, Heni Rachmawati. "The simple method of extra virgin olive oil nanoemulsion for oral consumption: the chance for enteral nanonutrition production", *International Journal of Biomedical Nanoscience and Nanotechnology*, 2020 10 words — < 1%
Crossref
-
- 24 jpsr.pharmainfo.in 10 words — < 1%
Internet
-
- 25 www.healthyfellow.com 10 words — < 1%
Internet
-
- 26 www.wjgnet.com 10 words — < 1%
Internet

27 Giuseppe Derosa, Angela D'Angelo, Pamela Maffioli. "The role of selected nutraceuticals in management of prediabetes and diabetes: An updated review of the literature", *Phytotherapy Research*, 2022

9 words — < 1%

Crossref

28 Javad Heshmati, Nazli Namazi, Mohammad-Reza Memarzadeh, Mohsen Taghizadeh, Fariba Kolahdooz. "Nigella sativa oil affects glucose metabolism and lipid concentrations in patients with type 2 diabetes: A randomized, double-blind, placebo-controlled trial", *Food Research International*, 2015

9 words — < 1%

Crossref

29 Lorenzo Flori, Lara Testai, Vincenzo Calderone. "The "irisin system": From biological roles to pharmacological and nutraceutical perspectives", *Life Sciences*, 2020

9 words — < 1%

Crossref

30 Md Ashrafal Alam, Riaz Uddin, Nusrat Subhan, Md Mahbubur Rahman, Preeti Jain, Hasan Mahmud Reza. "Beneficial Role of Bitter Melon Supplementation in Obesity and Related Complications in Metabolic Syndrome", *Journal of Lipids*, 2015

9 words — < 1%

Crossref

31 academic.oup.com

9 words — < 1%

Internet

32 core.ac.uk

9 words — < 1%

Internet

33 journals.lww.com

9 words — < 1%

Internet

34 lcm.amegroups.com

Internet

9 words — < 1%

35 [mdpi.com](https://www.mdpi.com)
Internet

9 words — < 1%

36 [turkiyeklinikleri.com](https://www.turkiyeklinikleri.com)
Internet

9 words — < 1%

37 [worldwidescience.org](https://www.worldwidescience.org)
Internet

9 words — < 1%

38 www.ajpsp.com
Internet

9 words — < 1%

39 www.tandfonline.com
Internet

9 words — < 1%

40 "Nutraceuticals and Natural Product Derivatives",
Wiley, 2019
Crossref

8 words — < 1%

41 Hira Choudhury, Manisha Pandey, Chua Kui Hua,
Cheah Shi Mun et al. "An update on natural
compounds in the remedy of diabetes mellitus: A systematic
review", Journal of Traditional and Complementary Medicine,
2018
Crossref

8 words — < 1%

42 L. WANG. "KEEPING QUALITY OF FRESH-CUT
BITTER GOURD (MOMORDICA CHARANTIA L.) AT
LOW TEMPERATURE OF STORAGE", Journal of Food Processing
and Preservation, 10/2007
Crossref

8 words — < 1%

43 Prapaipan Putthapiban, Weera
Sukhumthammarat, Chutintorn Sriphrapadang.

8 words — < 1%

"Concealed use of herbal and dietary supplements among Thai patients with type 2 diabetes mellitus", Journal of Diabetes & Metabolic Disorders, 2017

Crossref

44 brieflands.com 8 words — < 1%
Internet

45 www.thieme-connect.com 8 words — < 1%
Internet

46 Marcel Roberfroid, Glenn R. Gibson, Lesley Hoyles, Anne L. McCartney et al. "Prebiotic effects: metabolic and health benefits", British Journal of Nutrition, 2010 7 words — < 1%
Crossref

47 Remigio Usai, Stephen Majoni, Freeborn Rwere. "Natural products for the treatment and management of diabetes mellitus in Zimbabwe-a review", Frontiers in Pharmacology, 2022 7 words — < 1%
Crossref

48 hdl.handle.net 7 words — < 1%
Internet

49 "Prevention and Therapy of Type 2 Diabetes—What Is the Potential of Daily Water Intake and Its Mineral Nutrients?", Nutrients, 2017 6 words — < 1%
Crossref

50 Hasan Mihardja, Krisma Perdana Harja. "Acupuncture for Treating Hypertension and Type 2 Diabetes Mellitus as Comorbidities in Patients with COVID-19", Medical Acupuncture, 2021 6 words — < 1%
Crossref

51

Hongxiang Hui. "Hypoglycemic herbs and their action mechanisms", Chinese Medicine, 2009

Crossref

6 words — < 1%

EXCLUDE QUOTES ON

EXCLUDE BIBLIOGRAPHY ON

EXCLUDE SOURCES OFF

EXCLUDE MATCHES OFF