

Mechanical Ventilation Invasive

Setting: **Inpatient** Population: **NICU** Keywords: **ventilator associated pneumonia, device, VAP, ventilator, vent, breathing**

Last Updated: 03/18/2020. Copyright Elsevier BV. All rights reserved.

Clinical Description

Care of the hospitalized infant experiencing the need for controlled or assisted breathing through an artificial airway.

Key Information

- Artificial airways used in infants are generally cuffless because the narrowest portion of the cricoid cartilage acts as the cuff within smaller airways.
 - To reduce the risk of pulmonary aspiration, a swallow evaluation should be performed prior to oral intake or feeding.
 - Neonatal-sized and pediatric-sized tracheostomies should be changed on a regular schedule to prevent obstruction of the single lumen cannula.
 - Small airways and smaller diameter tubes are less compatible with speaking valves and tracheostomy plugs or caps due to increased resistance around the tube during exhalation. This increases the risk of airflow obstruction.
-

Clinical Goals

By transition of care

A. The patient will demonstrate achievement of the following goals:

- Effective Communication
- Optimal Device Function
- Absence of Device-Related Skin or Tissue Injury
- Absence of Ventilator-Induced Lung Injury

B. Patient, family or significant other will teach back or demonstrate education topics and points:

- Education: Overview
 - Education: Self Management
 - Education: When to Seek Medical Attention
-

Correlate Health Status

Correlate health status to:

- prenatal and birth history, comorbidity, congenital anomaly
 - gestational age, corrected age, day of life
 - sex
 - baseline assessment data
 - physiologic status
 - response to medication and interventions
 - barriers to accessing care and services
 - family/caregiver:
 - developmental level
 - health literacy
 - cultural and spiritual preferences
 - safety risks
 - social determinants of health
 - family interaction
 - plan for transition of care
-

Communication Impairment (Mechanical Ventilation, Invasive)

Signs/Symptoms/Presentation

- agitation
- artificial airway present inhibiting vocalization
- irritability
- soundless crying

Problem Intervention(s)

Ensure Effective Communication

- Acknowledge and validate intensity and complexity of infant's voicelessness and the inability of the parent/caregiver to hear the infant cry.
- Encourage seeing, touching, holding, skin-to-skin contact and response to infant cues, other than sound, to identify needs.
- Decrease parent/caregiver anxiety; provide honest and accurate information; encourage expression of feelings and participation in care.
- If longer-term airway, consider alternative communication methods that facilitate verbal sounds.

Associated Documentation

- Psychosocial Support
-

Device-Related Complication Risk (Mechanical Ventilation, Invasive)

Signs/Symptoms/Presentation

- air auscultated in stomach
- airflow out of mouth
- breath sounds unequal
- chest movement asymmetrical
- difficulty passing suction catheter
- excessive cough
- gastric distension
- gurgling sound from throat
- inability to ventilate
- mechanical ventilation with an artificial airway
- no airflow from device
- restlessness
- upper airway sounds increased
- work of breathing increased

Vital Signs

- heart rate increased
- respiratory rate increased
- SpO₂ (peripheral oxygen saturation) decreased
- TcCO₂/TcO₂ (transcutaneous carbon dioxide/oxygen) outside desired range

Laboratory Values

- PaCO₂ (arterial carbon dioxide) increased
- PaO₂ (partial pressure of arterial oxygen) decreased

Diagnostic Results

- CXR (chest x-ray) abnormal

Problem Intervention(s)

Optimize Device Care and Function

- Maintain head of bed elevation to minimize the risk of aspiration.
- Provide oral care regularly with subglottic suction to reduce the risk of infection; perform prior to tube manipulation.
- Assess tube size, depth, location and securement frequently to minimize the risk of tube displacement; regularly confirm placement with radiography or ultrasonography.
- Facilitate regular mechanical ventilator and humidification equipment checks to ensure proper function; monitor and manage ventilator and alarm settings.
- Provide humidification and evaluate need for suctioning to minimize risk of airway obstruction; regularly replace closed in-line suction equipment.
- Perform ongoing tracheostomy and stoma care to prevent infection; minimize excessive moisture around device ; replace tracheostomy regularly to prevent obstruction from secretions.
- Provide emergency equipment that includes appropriate-sized manual resuscitation bag, mask, suction equipment and cleaning supplies; replace device or assist breathing if displacement occurs.

Associated Documentation

- Airway Safety Measures
-

Skin and Tissue Injury (Mechanical Ventilation, Invasive)

Signs/Symptoms/Presentation

- bleeding
- hoarse cry
- laceration
- localized swelling
- redness
- skin blanching
- skin integrity disrupted
- stoma granulation
- stridor
- tracheal granuloma

Problem Intervention(s)

Maintain Skin and Tissue Health

- Monitor depth of suction catheter advancement to minimize the risk of internal tracheobronchial tissue injury.
- Reposition and resecure endotracheal tube regularly; ensure proper tube location.
- Monitor tightness of securement device, as well as skin and mucosal areas, regularly; consider skin barrier protection.
- Minimize pressure points and avoid traction on device; consider flexible extenders or props.
- Assess and monitor for the presence of bleeding that may indicate injury to tracheobronchial tissue. Notify provider for persistent bleeding.
- Anticipate the need for further treatment or procedure, if bleeding persists.
- Anticipate adjunct therapy, such as cool mist, racemic epinephrine, corticosteroid or heliox, for symptoms related to airway swelling or stridor after removal of tube.

Associated Documentation

- Device Skin Pressure Protection
-

Ventilator-Induced Lung Injury (Mechanical Ventilation, Invasive)

Signs/Symptoms/Presentation

- lung compliance decreasing
- oxygenation requirements increasing (e.g., FiO₂ or positive end expiratory pressure needs)
- ventilatory requirements increasing (e.g., minute volume, respiratory rate)

Vital Signs

- heart rate increased
- respiratory rate increased
- blood pressure increased or decreased
- SpO₂ (peripheral oxygen saturation) decreased
- TcCO₂/TcO₂ (transcutaneous carbon dioxide/oxygen) outside desired range

Laboratory Results

- oxygen index increased
- PaCO₂ (arterial carbon dioxide) increased
- PaO₂ (partial pressure of arterial oxygen) decreased

Diagnostic Results

- CXR (chest x-ray) abnormal

Problem Intervention(s)

Facilitate Lung-Protection Measures

- Provide oxygen therapy judiciously to maintain oxygenation goals; adjust to avoid hyperoxemia and oxygenation fluctuation.
- Maintain low inspiratory time to minimize barotrauma.
- Monitor and limit ventilator tidal volumes to minimize volutrauma; initiate low tidal-volume strategy (e.g., 4 to 6 mL/kg).
- Monitor and limit ventilator inspiratory pressure to reduce risk of barotrauma.
- Apply PEEP (positive end expiratory pressure) to minimize atelectasis; adjust for changes in lung compliance and oxygenation.
- Monitor fluid balance closely to minimize the risk of fluid overload.
- Monitor ventilator waveforms and promote patient-ventilator synchrony; adjust ventilator settings and sedation.

Prevent Ventilator-Associated Pneumonia

- Assess readiness to spontaneously breathe and extubate.
- Maintain head of bed elevation to minimize aspiration risk; consider lateral recumbent, reverse Trendelenburg and prone position.
- Provide ongoing oral care to reduce pathogens in oral cavity.
- Minimize ventilator circuit breaks; consider use of closed suction device.

Associated Documentation

- Lung Protection Measures

Associated Documentation

- Positioning, Body (Developmental Care)
- VAP Prevention Bundle

General Education

- admission, transition of care
- orientation to care setting, routine

- advance care planning
 - diagnostic tests/procedures
 - opioid medication management
 - oral health
 - medication management
 - pain assessment process
 - safe medication disposal
 - tobacco use, smoke exposure
 - treatment plan
-

Safety Education

- call light use
 - equipment/home supplies
 - fall prevention
 - harm prevention
 - infection prevention
 - MDRO (multidrug-resistant organism) care
 - personal health information
 - resources for support
-

Education: Overview

- description
 - indications
-

Education: Self-Management

- CPR education
 - VAP prevention
-

Education: When to Seek Medical Attention

- unresolved/worsening symptoms

References

- (2019). Walsh, B. K (Eds.), *Neonatal and pediatric respiratory care* . St. Louis: Elsevier. [Review Articles, Expert/Committee Opinion, Core Curriculum, Position Statements, Practice Bulletins]
- (2020). Cairo, J. M (Eds.), *Pilbeam's mechanical ventilation: Physiological and clinical applications* . St. Louis: Elsevier. [Review Articles, Expert/Committee Opinion, Core Curriculum, Position Statements, Practice Bulletins]
- American Association for Respiratory Care (AARC). Endotracheal suctioning of mechanically ventilated patients with artificial airways. *Respiratory Care*. 2010;55(6), 758-764. [Quality Measures, Clinical Practice Guidelines]
- American Association for Respiratory Care. Removal of the endotracheal tube - 2007 revision & update. *Respiratory Care*. 2007;52(1), 81-93. [Quality Measures, Clinical Practice Guidelines]
- Association of Women's Health, Obstetric and Neonatal Nurses (AWHONN). (2018). *Evidence-based clinical practice guideline: Neonatal skin care* . Washington, D. C.: Association of Women's Health, Obstetric and Neonatal Nurses. [Quality Measures, Clinical Practice Guidelines]
- Beck, J.; Emeriaud, G.; Liu, Y.; Sinderby, C. Neurally-adjusted ventilatory assist (NAVA) in children: A systematic review. *Minerva Anestesiologica*. 2016, August;82(8), 874-883. [Metasynthesis, Meta-analysis, Systematic Review]
- Bruschettini, M.; Zappettini, S.; Moja, L.; Calevo, M. G. Frequency of endotracheal suctioning for the prevention of respiratory morbidity in ventilated newborns. *Cochrane Database of Systematic Reviews*. 2016;(3) doi:10.1002/14651858.CD011493.pub2 [Metasynthesis, Meta-analysis, Systematic Review]
- College of Respiratory Therapists of Ontario. (2017). *Infection prevention & control clinical best practice guideline* . Source [Quality Measures, Clinical Practice Guidelines]
- Deuber, C.; Terhaar, M. Hyperoxia in very preterm infants: A systematic review of the literature. *Journal of Perinatal & Neonatal Nursing*. 2011;25(3), 268-274. doi:10.1097/JPN.0b013e318226ee2c [Metasynthesis, Meta-analysis, Systematic Review]
- EFCNI; Dubois, C.; Tissières, P.; Helder, O.; Mader, S.; Borghesi, A. (2018). *European Standards of Care for Newborn Health: Prevention of ventilator associated pneumonia* . Source .[Review Articles, Expert/Committee Opinion, Core Curriculum, Position Statements, Practice Bulletins]
- European Pressure Ulcer Advisory Panel; National Pressure Injury Advisory Panel; & Pan Pacific Pressure Injury Alliance. Prevention and treatment of pressure ulcers/injuries: Clinical practice guideline. *The International Guideline*. 2019;[Quality Measures, Clinical Practice Guidelines]
- Ferguson, K. N.; Roberts, C. T.; Manley, B. J.; Davis, P. G. Interventions to improve rates of successful extubation in preterm infants a systematic review and meta-analysis. *JAMA Pediatrics*. 2017;171(2), 165-174. doi:10.1001/jamapediatrics.2016.3015 [Metasynthesis, Meta-analysis, Systematic Review]

Gardner, S. L.; Carter, C. S.; Enzman-Hines, M.; Hernandez, J. A. (2016). *Merenstein and Gardner's handbook of neonatal intensive care*. St. Louis: Mosby, Elsevier. [Review Articles, Expert/Committee Opinion, Core Curriculum, Position Statements, Practice Bulletins]

Georgopoulos, D.; Prinianakis, G.; Kondili, E. Bedside waveforms interpretation as a tool to identify patient-ventilator asynchronies. *Intensive Care Medicine*. 2006;32(1), 34-47. doi:10.1007/s00134-005-2828-5L [Review Articles, Expert/Committee Opinion, Core Curriculum, Position Statements, Practice Bulletins]

Gillies, D.; Todd, D. A.; Foster, J. P.; Batuwitage, B. T. Heat and moisture exchangers versus heated humidifiers for mechanically ventilated adults and children. *Cochrane Database of Systematic Reviews*. 2017;(9) doi:10.1002/14651858.CD004711.pub3 [Metasynthesis, Meta-analysis, Systematic Review]

Greenough, A.; Rossor, T. E.; Sundaresan, A.; Murthy, V.; Milner, A. D. Synchronized mechanical ventilation for respiratory support in newborn infants. *Cochrane Database of Systematic Reviews*. 2016;(9) doi:10.1002/14651858.CD000456.pub5 [Metasynthesis, Meta-analysis, Systematic Review]

Hemsley, B.; Balandin, S. A metasynthesis of patient-provider communication in hospital for patients with severe communication disabilities: Informing new translational research. *Augmentative and Alternative Communication* (Baltimore, MD: 1985). 2014;30(4), 329-343. doi:10.3109/07434618.2014.955614 [Metasynthesis, Meta-analysis, Systematic Review]

Hockenberry, M. J.; Wilson, D.; Rodgers, C. C. (2019). *Wong's nursing care of infants and children*. St. Louis: Mosby, Elsevier. [Review Articles, Expert/Committee Opinion, Core Curriculum, Position Statements, Practice Bulletins]

Institute for Healthcare Improvement (IHI). (2012). *How-to Guide: Prevent Ventilator-Associated Pneumonia*. Source [Quality Measures, Clinical Practice Guidelines]

Kamlin, C. O. F.; Davis, P. G. Long versus short inspiratory times in neonates receiving mechanical ventilation. *Cochrane Database of Systematic Reviews*. 2003;(4) doi:10.1002/14651858.CD004503.pub2 [Metasynthesis, Meta-analysis, Systematic Review]

Klompas, M.; Branson, R.; Eichenwald, E. C.; Greene, L. R.; Howell, M. D.; Lee, G.; Magill, S. S.; Maragakis, L. L.; Priebe, G. P.; Speck, K.; Yoke, D. S.; Berenholtz, S. M. Strategies to prevent ventilator-associated pneumonia in acute care hospitals: 2014 update. *Infection Control & Hospital Epidemiology*. 2014;35(S2), S133-S154. doi:10.1017/S0899823X00193894 Source [Quality Measures, Clinical Practice Guidelines]

Kneyber, M. C.; De Luca, D.; Calderini, E.; Jarreau, P. H.; Javouhey, E.; Lopez-Herce, J.; et al. Recommendations for mechanical ventilation of critically ill children from the Paediatric Mechanical Ventilation Consensus Conference (PEMVECC). *Intensive Care Medicine*. 2017;43(12), 1764-1780. [Review Articles, Expert/Committee Opinion, Core Curriculum, Position Statements, Practice Bulletins]

Lai, M.; Inglis, G. D.; Hose, K.; Jardine, L. A.; Davies, M. W. Methods for securing endotracheal tubes in newborn infants. *Cochrane Database of Systematic Reviews*. 2014;(7) doi:10.1002/14651858.CD007805.pub2 [Metasynthesis, Meta-analysis, Systematic Review]

McCallion, N.; Davis, P. G.; Morley, C. J. Volume-targeted versus pressure-limited ventilation in the neonate. *Cochrane Database of Systematic Reviews*. 2005;(3) doi:10.1002/14651858.CD003666.pub2 [Metasynthesis, Meta-analysis, Systematic Review]

Mitchell, R. B.; Hussey, H. M.; Setzen, G.; Jacobs, I. N.; Nussenbaum, B.; Dawson, C.; et al. Clinical consensus statement: Tracheostomy care. *Otolaryngology-Head and Neck Surgery*. 2013;148(1), 6-20. doi:10.1177/

0194599812460376 [Review Articles, Expert/Committee Opinion, Core Curriculum, Position Statements, Practice Bulletins]

National Tracheostomy Safety Project. (2019). *Basic Care (child)*. Source [Review Articles, Expert/Committee Opinion, Core Curriculum, Position Statements, Practice Bulletins]

National Tracheostomy Safety Project. (2019). *Swallowing & Communication (child)*. Source [Review Articles, Expert/Committee Opinion, Core Curriculum, Position Statements, Practice Bulletins]

Peng, W.; Zhu, H.; Shi, H.; Liu, E. Volume-targeted ventilation is more suitable than pressure-limited ventilation for preterm infants: A systematic review and meta-analysis. *Archives of Disease in Childhood: Fetal and Neonatal Edition*. 2014;99(2), F158-165. doi:10.1136/archdischild-2013-304613 [Metasynthesis, Meta-analysis, Systematic Review]

Perrone, S.; Bracciali, C.; Di Virgilio, N.; Buonocore, G. Oxygen use in neonatal care: A two-edged sword. *Frontiers in Pediatrics*. 2017;4, 143. doi:10.3389/fped.2016.00143 [Review Articles, Expert/Committee Opinion, Core Curriculum, Position Statements, Practice Bulletins]

Peterson-Carmichael, S.; Seddon, P. C.; Cheifetz, I. M.; Frerichs, I.; Hall, G. L.; Hammer, J.; et al. An official American Thoracic Society/European Respiratory Society Workshop Report: Evaluation of respiratory mechanics and function in the pediatric and neonatal intensive care units. *Annals of the American Thoracic Society*. 2016;13(2), S1-S11. [Review Articles, Expert/Committee Opinion, Core Curriculum, Position Statements, Practice Bulletins]

Registered Nurses' Association of Ontario (RNAO). (2002). *Establishing therapeutic relationships*. Toronto, Canada: Registered Nurses Association of Ontario (RNAO). [Quality Measures, Clinical Practice Guidelines]

Registered Nurses' Association of Ontario. (2015). *Person- and family-centred care*. Source [Quality Measures, Clinical Practice Guidelines]

Restrepo, R. D.; Walsh, B. K. Humidification during invasive and noninvasive mechanical ventilation: 2012. *Respiratory Care*. 2012;57(5), 782-788. [Quality Measures, Clinical Practice Guidelines]

Rivas-Fernandez, M.; Roque I Figuls, M.; Diez-Izquierdo, A.; Escribano, J.; Balaguer, A. Infant position in neonates receiving mechanical ventilation. *Cochrane Database of Systematic Reviews*. 2016;(11) doi:10.1002/14651858.CD003668.pub4 [Metasynthesis, Meta-analysis, Systematic Review]

Schmölzer, G. M.; Roehr, C. C. Techniques to ascertain correct endotracheal tube placement in neonates. *Cochrane Database of Systematic Reviews*. 2014;(9) doi:10.1002/14651858.CD010221.pub2 [Metasynthesis, Meta-analysis, Systematic Review]

Sweet, D. G.; Carnielli, V.; Greisen, G.; Hallman, M.; Ozek, E.; te Pas, A.; . . . & Speer, C. P. European consensus guidelines on the management of respiratory distress syndrome-2019 update. *Neonatology*. 2019;115(4), 432-450. [Quality Measures, Clinical Practice Guidelines]

Tan, B.; Zhang, F.; Zhang, X.; Huang, Y. L.; Gao, Y. S.; Liu, X.; Li, Y. L.; Qiu, J. F. Risk factors for ventilator-associated pneumonia in the neonatal intensive care unit: A meta-analysis of observational studies. *European Journal of Pediatrics*. 2014;173(4), 427-434. doi:10.1007/s00431-014-2278-6 [Review Articles, Expert/Committee Opinion, Core Curriculum, Position Statements, Practice Bulletins]

Taylor, J. E.; Hawley, G.; Flenady, V.; Woodgate, P. G. Tracheal suctioning without disconnection in intubated ventilated neonates. *Cochrane Database of Systematic Reviews*. 2011;(12) doi:10.1002/14651858.CD003065.pub2 [Metasynthesis, Meta-analysis, Systematic Review]

Wallis, C.; Alexopoulou, E.; Antón-Pacheco, J. L.; et al. ERS statement on tracheomalacia and bronchomalacia in children. *European Respiratory Journal*. 2019;54, 1900382. doi:10.1183/13993003.00382-2019 Source [Quality Measures, Clinical Practice Guidelines]

Walsh, B. K.; Crotwell, D. N.; Restrepo, R. D. Capnography/Capnometry during mechanical ventilation: 2011. *Respiratory Care*. 2011;56(4), 503-509. [Quality Measures, Clinical Practice Guidelines]

Weiner, G. M. (2016). *Textbook of neonatal resuscitation*. Elk Grove Village: American Academy of Pediatrics. [Metasynthesis, Meta-analysis, Systematic Review]

Wheeler, K. I.; Klinenberg, C.; Morley, C. J.; Davis, P. G. Volume-targeted versus pressure-limited ventilation for preterm infants: A systematic review and meta-analysis. *Neonatology*. 2011;100(3), 219-227. doi:10.1159/000326080 [Metasynthesis, Meta-analysis, Systematic Review]

de França, D. F.; do Nascimento Cassiano, A.; Vitorino, A. B. F.; de Oliveira, S. I. M.; e Silva, R. K. C.; Souza, J. R. S.; de Souza, N. L. Bundle method in reducing pneumonia associated with mechanical ventilation in newborns and children: Integrative review. *International Archives of Medicine*. 2016;9(1)[Review Articles, Expert/Committee Opinion, Core Curriculum, Position Statements, Practice Bulletins]
