



COVID-19 Drug Therapy

Tim Smith, PharmD, BCPS; Jennifer Bushek, PharmD; Aimée LeClaire, PharmD, BCPS;
Tony Prosser, PharmD

Clinical Drug Information | Clinical Solutions

What's been updated:

- Remdesivir
 - Recommendation for use from NIH COVID-19 treatment guidelines(133)
 - New randomized, double-blind, placebo-controlled, multicenter trial (150)
 - Safety concerns addition
- Chloroquine
 - Recommendation against the use of high-dose chloroquine from NIH COVID-19 treatment guidelines (133)

Highlights:

- There are no specific therapies approved by the U.S. Food and Drug Administration (FDA) for severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), the virus that causes coronavirus disease 2019 COVID-19. Several agents are being used under clinical trial and compassionate use protocols based on *in vitro* activity (against SARS-CoV-2 or related viruses) and on limited clinical experience. Efficacy has not been established for any drug therapy.
- **Antimicrobials with potential activity against SARS-CoV-2:**
 - Remdesivir – Investigational antiviral available under an FDA Emergency Use Authorization (EUA); several large clinical trials are underway with preliminary data suggesting clinical benefit.
 - Chloroquine – *In vitro* and limited clinical data suggest potential benefit; FDA EUA issued to facilitate availability.
 - Hydroxychloroquine – *In vitro* and limited clinical data suggest potential benefit; FDA EUA issued to facilitate availability.
 - Lopinavir; Ritonavir – Role in the treatment of COVID-19 is unclear. Preclinical data suggested potential benefit; however, more recent data has failed to confirm.
 - Favipiravir – Investigational use is being studied.

- **Adjunctive / supportive care:**

- Anticoagulation – Venous thromboembolism prophylaxis with low molecular weight heparin (LMWH) recommended for all hospitalized patients.
- Azithromycin – Used in some protocols based on theoretical mechanism and limited preliminary data as adjunct therapy.
- Bronchodilators – No routine role for inhaled bronchodilators in the management of COVID-19; metered-dose inhalers (MDI) preferred over nebulized therapy due to the risk of viral transmission.
- Corticosteroids – Not recommended for viral pneumonia; use may be considered for patients with refractory shock or acute respiratory distress syndrome.
- COVID-19 convalescent plasma – Investigational use is being studied.
- Fibrinolytics – Severe COVID-19 infection is associated with coagulopathy; fibrin deposition in the pulmonary microvasculature is a causative factor in the development of ARDS.
- Immunomodulating agents [Interleukin Receptor Antagonists, Janus Kinase (JAK) inhibitors] – Used in some protocols based on theoretical mechanisms and limited preliminary data as adjunct therapy.
- Inhaled pulmonary vasodilators – No evidence for routine in acute respiratory failure; use may be considered in specific patients with acute respiratory distress syndrome (ARDS) as a temporizing measure.
- NSAIDs – The FDA continues to investigate the use of NSAIDs; concern for potential worsening of COVID-19 symptoms has been suggested, but confirmatory clinical data is lacking.

According to the World Health Organization (WHO), the Centers for Disease Control and Prevention (CDC), the FDA, and the National Institutes of Health (NIH), there are currently no medications or vaccines proven to be effective for the treatment or prevention of SARS-CoV-2. (1) (2) (3) (133)

Generally, pharmacologic treatment is not recommended for young, healthy patients with mild symptoms and no underlying comorbid conditions.(12) (13)

Antimicrobials with potential activity against SARS-CoV-2:

Remdesivir (GS-5734):

- Classification: Investigational Nucleoside Analogue
- Rationale for Use: Remdesivir is a broad-spectrum antiviral with in vitro activity against coronaviruses.(10) (14) (38) (39) (41) (42) (43) (44)
- Mechanism of Action: Remdesivir is a monophosphoramidate prodrug of remdesivir-triphosphate (RDV-TP), an adenosine analog that acts as an inhibitor of RNA-dependent RNA polymerases (RdRps). Remdesivir-TP competes with adenosine-triphosphate for incorporation into nascent viral RNA chains. Once incorporated into the viral RNA at position i, RDV-TP

terminates RNA synthesis at position i+3. Because RDV-TP does not cause immediate chain termination (i.e., 3 additional nucleotides are incorporated after RDV-TP), the drug appears to evade proofreading by viral exoribonuclease (an enzyme thought to excise nucleotide analog inhibitors).(10) (14) (38) (39) (41) (42) (43) (44) (45)

- FDA Emergency Use Authorization (EUA): (146)
 - Remdesivir is not an FDA-approved medication.
 - The EUA allows for the emergency use of remdesivir to treat suspected or laboratory confirmed COVID-19 in adults and children hospitalized with severe disease.
 - The EUA defines severe COVID-19 disease as patients with an oxygen saturation (SpO₂) of 94% or less on room air, or requiring supplemental oxygen, or requiring mechanical ventilation or extracorporeal membrane oxygenation (ECMO).
- Evidence / Experience:
 - The NIH COVID-19 treatment guidelines recommend the use of remdesivir in hospitalized patients with severe COVID-19; however, its use in patients with mild or moderate COVID-19 is not recommended outside of clinical trials.(133)
 - Preliminary data from a Phase III trial [i.e., Adaptive COVID-19 Treatment Trial (ACTT)] suggest remdesivir may provide clinical benefit in the treatment of hospitalized patients with advanced COVID-19 and lung involvement.(145) (146)
 - The median time to recovery was 11 days in the remdesivir-treated group compared to 15 days in placebo group (hazard ratio: 1.31; 95% CI: 1.12 to 1.54, p < 0.001).
 - Mortality was 8% for the remdesivir group vs. 11.6% for the placebo group (p = 0.059)
 - A randomized, double-blind, placebo-controlled, multicenter trial evaluating efficacy and safety of remdesivir in SARS-CoV-2 infected hospitalized adults (concurrent treatment with corticosteroids, interferons, and lopinavir; ritonavir was permitted).(150)
 - No difference between remdesivir (n = 158) and placebo (n = 79) in the time to clinical improvement (median, 21 vs. 23 days; hazard ratio, 1.23; 95% CI: 0.87 to 1.75). 28-day mortality was similar between remdesivir (14%) and placebo (13%) (95% CI: -8.1 to 10.3).
 - Adverse events reported in 66% of remdesivir patients and 64% of placebo patients. Remdesivir was stopped early in 12% of patients due to adverse events.
 - Preliminary data from open-label compassionate use in patients with severe disease was analyzed.(124)
 - Clinical improvement, as defined by improvement in oxygen support, was reported in 36 of 53 patients (68%).
 - Clinical improvement was less frequent in those receiving invasive ventilation and in patients 70 years or older.
 - Several clinical trials evaluating the efficacy of remdesivir in patients infected with SARS-CoV-2 are currently being conducted.(132)
- Safety Concerns: (133) (146)
 - Caution in patients with renal impairment due to formulation with sulfobutyl ether beta-cyclodextrin sodium (SBECD)
 - Infusion-related reactions
 - Risk for elevated hepatic enzymes

Chloroquine:

- Classification: Antimalarial
- Rationale for Use: Chloroquine has *in vitro* activity against SARS-CoV-2 and may have immunomodulating properties.(13) (14) (15) (17)
- Mechanism of Action: Mechanisms may include inhibition of viral enzymes or processes such as viral DNA and RNA polymerase, viral protein glycosylation, virus assembly, new virus particle transport, and virus release. Other mechanisms may also involve ACE2 cellular receptor inhibition, acidification at the surface of the cell membrane inhibiting fusion of the virus, and immunomodulation of cytokine release.(14) (15) (29) (30) (31) (32) (33)
- FDA Emergency Use Authorization (EUA): (66) (67)
 - Chloroquine is not FDA-approved for the treatment of COVID-19.
 - EUA is to facilitate the availability of chloroquine during the COVID-19 pandemic to treat patients for whom a clinical trial is not available, or participation is not feasible.
 - The EUA states treatment is for adult and adolescent patients weighing 50 kg or more who are hospitalized with COVID-19.
 - Authorized chloroquine is limited to product supplied from the Strategic National Stockpile (SNS) and will be distributed to authorized health care systems and providers.
- Evidence / Experience:
 - Due to a lack of clinical data, the NIH COVID-19 treatment guidelines do not give recommendations for or against the use of chloroquine; however, if used, guidelines advise monitoring for adverse events, including QT interval prolongation. The NIH recommends against the use of high-dose, twice-daily chloroquine due to a higher risk of toxicities.(133)
 - Pre-clinical data *in vitro* suggest chloroquine has activity against SARS-CoV-2.(13) (14) (15)
 - There have been reports of potential benefit in inhibiting the exacerbation of pneumonia patients with SARS-CoV-2 infection; however, specific data are not available.(13)
 - Some protocols include recommendations for use.(12) (21) (22)
 - Additional data regarding clinical efficacy for COVID-19 are being evaluated.(16) (31)
- Safety Concerns: (46) (49)
 - Use in COVID-19 patients outside of clinical trials or in a nonhospital setting is not recommended due to the potential for serious adverse events and drug interactions.(141)
 - Risk of cardiac arrhythmias (e.g., QT prolongation)
 - Seriously ill patients with comorbidities have an increased risk of serious arrhythmias.(125)
 - Avoid other QT prolonging agents whenever feasible.(125)
 - In a parallel, double-masked, randomized clinical trial (n = 81), two doses of chloroquine were analyzed (600 mg PO twice daily for 10 days vs. 450 mg PO twice daily on day 1 then 450 mg PO once daily for 4 days).(126)
 - The high-dose group had a higher incidence of QT interval greater than 500 milliseconds (18.9%) compared with the low-dose group (11.1%); therefore, the study was unmasked and all patients were reverted to the low-dose group.
 - Risk of retinal damage, especially with long term use

- Caution in patients with G6PD deficiency
- Caution in diabetics
- Significant drug interactions

Hydroxychloroquine:

- Classification: Antimalarial
- Rationale for Use: Hydroxychloroquine has *in vitro* activity against SARS-CoV-2 and may have immunomodulating properties.(13) (14) (15) (17)
- Mechanism of Action: Mechanisms may include inhibition of viral enzymes or processes such as viral DNA and RNA polymerase, viral protein glycosylation, virus assembly, new virus particle transport, and virus release. Other mechanisms may also involve ACE2 cellular receptor inhibition, acidification at the surface of the cell membrane inhibiting fusion of the virus, and immunomodulation of cytokine release.(14) (15) (29) (30) (31) (32) (33)
- FDA Emergency Use Authorization (EUA): (66) (68)
 - Hydroxychloroquine is not FDA-approved for the treatment of COVID-19.
 - EUA is to facilitate the availability of hydroxychloroquine during the COVID-19 pandemic to treat patients for whom a clinical trial is not available, or participation is not feasible.
 - The EUA states treatment is for adult and adolescent patients weighing 50 kg or more who are hospitalized with COVID-19.
 - Authorized hydroxychloroquine is limited to product supplied from the Strategic National Stockpile (SNS) and will be distributed to authorized health care systems and providers.
- Evidence / Experience:
 - Due to a lack of clinical data, the NIH COVID-19 treatment guidelines do not give recommendations for or against the use of hydroxychloroquine; however, if used, guidelines advise monitoring for adverse events including QT interval prolongation (133)
 - Pre-clinical *in vitro* data suggest hydroxychloroquine has activity against SARS-CoV-2.(12) (15) (17) (18) (21)
 - One *in vitro* study suggests that hydroxychloroquine may be more potent than chloroquine.(15)
 - Hydroxychloroquine exhibited a higher *in vitro* antiviral effect compared to chloroquine when drug was added prior to the viral challenge.
 - The EC50 values for chloroquine were greater than 100 microM at 24 hours and 18.01 microM at 48 hours.
 - The EC50 values for hydroxychloroquine were 6.25 microM at 24 hours and 5.85 microM at 48 hours.
 - An open-label, non-randomized clinical trial compared hydroxychloroquine treatment (n = 26) to an untreated negative control group.(27)
 - Preliminary data showed the proportion of patients that had negative PCR results significantly differed between treated patients and untreated controls.
 - On day 6, 70% of hydroxychloroquine-treated patients were virologically cured compared to 12.5% in the untreated control group.

- A parallel-group, randomized trial (n = 62) of hospitalized patients with non-severe COVID-19 compared 5 days of hydroxychloroquine to standard treatment.(69)
 - Fever recovery time was shortened in the hydroxychloroquine group (2.2 days) compared to standard therapy (3.2 days).
 - Cough recovery time was shortened in the hydroxychloroquine group (2 days) compared to standard therapy (3.1 days).
- A prospective review assessed virologic and clinical outcomes of 11 hospitalized patients who received hydroxychloroquine and azithromycin.(88)
 - Within 5 days, 1 patient died, 2 were transferred to the ICU, and 1 patient had therapy discontinued due to QT prolongation.
 - Nasopharyngeal swabs were still positive for SARS-CoV-2 in 8 of 10 patients 5 to 6 days after treatment initiation.
- In a preliminary report of a multicenter, parallel, open-label, randomized trial in 150 adult hospitalized patients, hydroxychloroquine (n = 75) was added to standard therapy.(127)
 - The overall 28-day negative viral conversion rate was not different between the two groups (85.4% hydroxychloroquine vs 81.3% control, p = 0.341).
 - Negative conversion rates on days 4, 7, 10, and 21 were also similar between the groups.
 - There was no difference in the 28-day symptom alleviation rate; however, a significant efficacy on symptom alleviation was observed when the confounding effects of antiviral agents were removed in the post-hoc analysis.
 - Additionally, there was a greater reduction in C-reactive protein (CRP) (6.986 mg/L hydroxychloroquine vs 2.723 mg/L control, p = 0.045) and a nonsignificant increase in recovery of lymphopenia in the hydroxychloroquine group.
 - Adverse events were reported in 8.8% of the control group and 30% of the hydroxychloroquine group.
- Preliminary data from a retrospective analysis (n = 368) assessed the use of either hydroxychloroquine (n = 97) or hydroxychloroquine plus azithromycin (n = 133) in addition to standard of care compared to standard of care alone (n = 158).(134)
 - The cohort consisted of only men in the Veterans Health Administration medical centers with a median age older than 65 years, the majority of whom were black.
 - The primary outcomes were death and the need for mechanical ventilation.
 - Rates of death in the hydroxychloroquine group, the hydroxychloroquine plus azithromycin group, and the standard treatment group were 27.8%, 22.1%, and 11.4%, respectively. Compared to standard therapy, the risk of death from any cause was higher in the hydroxychloroquine group (adjusted HR 2.61; 95% CI, 1.10 to 6.17; p = 0.03), but not in the hydroxychloroquine plus azithromycin group (adjusted HR 1.14; 95% CI, 0.56 to 2.32; p = 0.72).
 - Rates of ventilation in the hydroxychloroquine group, the hydroxychloroquine plus azithromycin group, and the standard treatment group were 13.3%, 6.9%, and 14.1%, respectively. Compared to standard therapy, the risk of ventilation was similar in both the hydroxychloroquine group (adjusted HR 1.43; 95% CI, 0.53 to 3.79; p =

- 0.48) and the hydroxychloroquine plus azithromycin group (adjusted HR 0.43; 95% CI, 0.16 to 1.12; p = 0.09).
- An observational trial (n = 1,376) examined the association between hydroxychloroquine use and intubation or death at a large medical center.(149)
 - Hydroxychloroquine was not associated with a significantly higher or lower risk of intubation or death (HR 1.04; 95% CI, 0.82 to 1.32); similar results were noted when adjusted for propensity score.
 - Hydroxychloroquine-treated patients were more severely ill at baseline.
 - Due to wide confidence intervals and the observational nature of the trial, the authors stated that the results should not be utilized to rule out either benefit or harm of hydroxychloroquine and suggested further randomized clinical trials to test efficacy.
 - An observational study of 1,438 hospitalized patients assessed mortality in patients receiving hydroxychloroquine (n = 271), azithromycin (n = 211), or both (n = 735) compared to patients who received neither of these agents (n = 221).(155)
 - There was no difference in mortality in patients treated with hydroxychloroquine (HR 1.08; 95% CI, 0.63 to 1.85), azithromycin (HR 0.56; 95% CI, 0.26 to 1.21), or both (HR 1.35; 95% CI, 0.76 to 2.4) compared with no use of these agents.
 - In logistic models, cardiac arrest was significantly more likely in patients receiving hydroxychloroquine plus azithromycin (OR 2.13; 95% CI, 1.12 to 4.05) compared to patients receiving neither drug; however, this was not the case in patients receiving either drug alone.
 - In adjusted logistic regression models, there were no significant differences in the relative likelihood of abnormal electrocardiogram findings.
 - Patient receiving hydroxychloroquine with or without azithromycin were overall sicker on presentation.
 - Some protocols have recommendations for use.(12) (21)
 - Additional data regarding clinical efficacy for COVID-19 are being evaluated.(16) (31)
 - Safety Concerns: (47) (49)
 - Use in COVID-19 patients outside of clinical trials or in a nonhospital setting is not recommended due to the potential for serious adverse events and drug interactions.(141)
 - Risk of cardiac arrhythmias (e.g., QT prolongation)
 - Seriously ill patients with comorbidities have an increased risk of serious arrhythmias.(125)
 - Avoid other QT prolonging agents whenever feasible.(125)
 - Risk of retinal damage, especially with long term use
 - Caution in patients with G6PD deficiency
 - Caution in diabetics
 - Significant drug interactions

Lopinavir; Ritonavir:

- Classification: HIV Protease Inhibitor
- Rationale for Use: *In vitro* and animal model studies show potential activity for other coronaviruses (SARS-CoV and MERS-CoV).(4) (52) (53) (54)
- Mechanism of Action: Lopinavir and ritonavir may bind to M^{pro}, a key enzyme for coronavirus replication. This may suppress coronavirus activity.(55)
- Evidence / Experience:
 - Due to unfavorable pharmacodynamics and negative clinical trial data, the NIH COVID-19 treatment guidelines recommend against the use of lopinavir; ritonavir or other HIV protease inhibitors outside of clinical trials.(133) Similarly, ESICM and SCCM Surviving Sepsis Campaign recommendations suggest against the routine use of lopinavir; ritonavir in critically ill adults with COVID-19.(26)
 - Pre-clinical data show activity for other coronaviruses.(4) (52) (53) (54)
 - A randomized, controlled, open-label trial involving hospitalized patients with confirmed SARS-CoV-2 infection (n = 199), analyzed treatment with lopinavir; ritonavir.(23)
 - Not associated with a difference from standard of care in the time to clinical improvement (median, 16 days versus 16 days; hazard ratio 1.31; 95% CI, 0.95 to 1.80; p = 0.09); percentages of patients with detectable viral RNA were similar; 28 day mortality was also similar (19.2% vs. 25%, respectively).
 - A retrospective cohort study of hospitalized patients reviewing clinical course and risk factors for mortality included 29 patients who received lopinavir; ritonavir.(24)
 - No difference was noted in the duration of viral shedding after treatment with lopinavir; ritonavir.
- Safety Concerns: (45) (49)
 - Risk of cardiac arrhythmias (e.g., QT prolongation)
 - Seriously ill patients with comorbidities have an increased risk of serious arrhythmias.(125)
 - Avoid other QT prolonging agents whenever feasible.(125)
 - Caution in patients with hepatic disease or hepatitis
 - Significant drug interactions

Favipiravir:

- Classification: Investigational RNA-Dependent RNA Polymerase Inhibitor
- Rationale for Use: Favipiravir is a broad-spectrum antiviral with *in vitro* activity against RNA viruses.(14) (18) (75)
- Mechanism of Action: Favipiravir is an RNA-dependent RNA polymerase (RdRp) inhibitor that inhibits viral RNA synthesis.(14) (18) (75)
- Evidence / Experience:
 - In a non-randomized, controlled, open-label trial, the efficacy of favipiravir (n = 35) in treating patients with laboratory confirmed COVID-19 was compared against lopinavir; ritonavir (n = 45); both treatments were given in combination with inhaled interferon alpha.(142)

- Time to viral clearance was shorter for favipiravir (median, 4 days; range, 2.5 to 9 days) than for lopinavir; ritonavir (median, 11 days; range 8 to 13 days; $p < 0.001$).
 - Chest imaging improvement rate at treatment day 14 was 91% for favipiravir vs. 62% for lopinavir; ritonavir ($p = 0.004$)
 - A prospective, randomized, controlled, open-label multicenter trial compared favipiravir ($n = 116$) against umifenovir [Arbidol] ($n = 120$) in treating patients with COVID-19; both treatments were given in combination with conventional therapy.(143)
 - No difference in clinical recovery rate at treatment Day 7 (61% for favipiravir vs. 52% for umifenovir; $p = 0.1396$; 95% CI: -0.0305 to 0.2213).
 - Additional data regarding clinical efficacy for COVID-19 are being evaluated.(132)
- **Safety Concerns:**
 - Contraindicated in pregnancy due to early embryonic death and teratogenicity observed in animal studies.

Adjunctive/Supportive therapy:

Anticoagulation:

- Venous thromboembolism (VTE) prophylaxis with LMWH is recommended for all hospitalized patients with COVID-19 infection. Withhold VTE prophylaxis for active bleeding, platelet count less than $25 \times 10^9/L$, or fibrinogen less than 0.5 g/L.(80) (81) (91) (95) (102)
 - Use fondaparinux in patients with a history of heparin-induced thrombocytopenia.(95)
 - Use unfractionated heparin or reduced-dose LMWH in patients with creatinine clearance less than 30 mL/minute.(103)
 - Use mechanical thromboprophylaxis in patients where anticoagulants are contraindicated or unavailable.(95)
 - Data are insufficient to recommend for or against the use of increasing anticoagulant doses for VTE prophylaxis in hospitalized COVID-19 patients outside of a clinical trial.(133)
 - Hospitalized patients with COVID-19 should not routinely be discharged on VTE prophylaxis. Weighing the individual patient's VTE risk factors and bleeding risk in addition to feasibility, consider extended thromboprophylaxis with a regulatory-approved regimen (e.g., betrixaban, rivaroxaban) after discharge.(95) (133)
 - Aspirin could be considered for COVID-19 VTE prophylaxis if criteria for post-discharge thromboprophylaxis are met.(95)
 - Elevated D-dimer has been noted in COVID-19 patients requiring hospitalization and has been associated with increased mortality. Limited data suggest a decrease in mortality in patients with severe COVID-19 infection or markedly elevated D-dimer concentrations (more than 6 times the upper limit of normal) who were given LMWH or heparin VTE prophylaxis.(80) (81) (91) (102)
- Therapeutic-intensity anticoagulation is not recommended in the management of COVID-19 in the absence of confirmed or suspected VTE.(95) (128)

- In patients already anticoagulated for VTE or atrial fibrillation, continue therapeutic anticoagulation. Consider withholding therapeutic anticoagulation in these patients for platelet count less than $50 \times 10^9/L$ or fibrinogen less than 1 g/L.(102)
- Treat all COVID-19 patients with confirmed or suspected VTE with therapeutic anticoagulation for at least 3 months. Therapeutic anticoagulation may be discontinued at 3 months if the patient has recovered from COVID-19 and has no ongoing risk factors for thrombosis or other indications for anticoagulation.(128)
- Increasing the intensity of anticoagulation (i.e., from standard-intensity prophylaxis to intermediate-intensity prophylaxis or from intermediate-intensity prophylaxis to therapeutic-intensity prophylaxis) may be reasonable in COVID-19 patients who experience recurrent clotting of access devices or extracorporeal circuits.(95)

Azithromycin:

- Classification: Macrolide Antibacterial
- Rationale for Use: Azithromycin may prevent bacterial superinfection, and macrolides may have immunomodulatory properties to work as adjunct therapy.(27) (34) (35) (36) (37)
- Mechanism of Action: Macrolides may have immunomodulatory properties in pulmonary inflammatory disorders. They may downregulate inflammatory responses and reduce the excessive cytokine production associated with respiratory viral infections; however, their direct effects on viral clearance are uncertain. Immunomodulatory mechanisms may include reducing chemotaxis of neutrophils (PMNs) to the lungs by inhibiting cytokines (i.e., IL-8), inhibition of mucus hypersecretion, decreased production of reactive oxygen species, accelerating neutrophil apoptosis, and blocking the activation of nuclear transcription factors.(34) (35) (36) (37)
- Evidence / Experience:
 - Due to the potential for toxicities, the NIH COVID-19 treatment guidelines recommend against the use of azithromycin in combination with hydroxychloroquine outside of clinical trials.(133)
 - In an open-label, non-randomized clinical trial of hydroxychloroquine (n = 26), azithromycin was administered in combination with hydroxychloroquine to prevent bacterial superinfection in 6 patients.(27)
 - Preliminary data suggest the potential for benefit as adjunct therapy.
 - On day 6, all patients treated with the combination (hydroxychloroquine and azithromycin) were virologically cured compared to 57.1% of patients treated with hydroxychloroquine alone (n= 20).
 - In a retrospective analysis of a multicenter cohort study (n = 349) in patients with MERS-CoV, 136 patients received macrolide therapy in combination with antiviral treatment.(28)
 - Macrolide therapy was not associated with a reduction in 90-day mortality compared to the control group (adjusted OR: 0.84; 95% CI: 0.47 to 1.51; p = 0.56).
 - Sensitivity analysis excluding patients who received macrolides after day 3 showed similar results (adjusted OR: 0.7; 95% CI: 0.39 to 1.28; p = 0.25).
 - A prospective review assessed virologic and clinical outcomes of 11 hospitalized patients who received hydroxychloroquine and azithromycin.(88)

- Within 5 days, 1 patient died, 2 were transferred to the ICU, and 1 patient had therapy discontinued due to QT prolongation.
 - Nasopharyngeal swabs were still positive for SARS-CoV-2 in 8 of 10 patients 5 to 6 days after treatment initiation.
- Safety Concerns: (48) (49)
 - Risk of cardiac arrhythmias (e.g., QT prolongation)
 - Seriously ill patients with comorbidities have an increased risk of serious arrhythmias.(125)
 - Avoid other QT prolonging agents whenever feasible.(125)
 - Significant drug interactions

Bronchodilators

- Most patients with COVID-19 do not need inhaled bronchodilator therapy. There is no role for inhaled bronchodilators in the management of COVID-19 unless the patient has underlying asthma or chronic obstructive pulmonary disease (COPD).(57) (61)
 - MDIs are preferred due to the potential for generation of aerosols that may increase the risk of viral transmission with nebulized therapy.(57) (61)
 - Due to concerns about supply chain interruption, some institutions are developing an MDI canister reassignment protocol to address potential shortages. An MDI canister reassignment protocol should emphasize hand hygiene and dual canister disinfection and avoid inadvertent sources of transmission.(57) (79)

Corticosteroids:

- Corticosteroid therapy is not recommended for viral pneumonia; however, use may be considered for patients with refractory shock or ARDS.(1) (7) (26) (62) (63) (64) (133)

COVID-19 Convalescent Plasma: (22)

- Classification: Plasma collected from persons who have recovered from COVID-19 that may contain antibodies to SARS-CoV-2
- Rationale for Use: Clinical trials are being conducted to evaluate the use of COVID-19 convalescent plasma to treat patients with severe or immediately life-threatening COVID-19 infections. COVID-19 convalescent plasma is not intended for prevention of the infection.
 - To participate in these trials, investigators should submit requests to the FDA for investigational use under the traditional IND regulatory pathway.
 - In addition to clinical trials, licensed physicians may obtain COVID-19 convalescent plasma for an individual patient through the process of single patient eINDs.
- Evidence / Experience:
 - Due to a lack of clinical data, the NIH COVID-19 treatment guidelines do not give recommendations for or against the use of convalescent plasma.(133)
 - In a case series of 5 critically ill patients with confirmed COVID-19 and ARDS, patients received convalescent plasma.(65)

- Treatment: 2 consecutive transfusions of 200 mL to 250 mL of convalescent plasma (total dose: 400 mL) with a SARS-CoV-2-specific antibody (IgG) titer greater than 1:1,000 on the same day it was obtained from the donor.
- Patient criteria included:
 - Severe pneumonia with rapid progression and continuously high viral load despite antiviral treatment
 - PAO_2/FIO_2 less than 300
 - Mechanical ventilation
- After plasma infusion, body temperature normalized within 3 days in 4 of 5 patients, Sequential Organ Failure Assess (SOFA) score decreased and PAO_2/FIO_2 increased within 12 days.
- Viral loads decreased and became negative within 12 days after the transfusion with the SARS-CoV-2-specific ELISA and neutralizing antibody titers increased after the transfusion.
- ARDS resolved in 4 patients by day 12 after the transfusion and 3 patients were weaned from mechanical ventilation within 2 weeks of treatment.
- In a case series of 6 critically ill patients with confirmed COVID-19 with abnormalities on chest CT (with the exception of 1 patient) who were deteriorating while receiving standard treatment, patients received convalescent plasma.(135)
 - Patients received at least 1 cycle (range, 1 to 3 cycles) of convalescent plasma (200 mL per cycle) over 30 minutes.
 - All patients had improved symptoms and chest CT and were discharged from the hospital.

Fibrinolytics:

- Rationale for Use: Severe COVID-19 infection is associated with coagulopathy, specifically a prothrombotic disseminated intravascular coagulation (DIC) with high rates of VTE, vascular occlusive events, and central line thrombosis and pulmonary congestion with microvascular thrombosis and occlusion observed on pathology. Fibrin deposition in the pulmonary microvasculature is a causative factor in the development of ARDS.(102) (136)
- Mechanism of Action: Fibrinolytic agents act by converting plasminogen to plasmin on the surface of existing thrombi, thereby initiating local fibrinolysis.(137)
- Evidence / Experience:
 - Alteplase:
 - In a case series of 3 critically ill patients with ARDS and respiratory failure, patients received alteplase followed by intravenous heparin.(136)
 - PaO_2/FiO_2 (P/F) ratio improved by 11% to 100% in all 3 patients; however, the improvements were transient.
 - Additional data regarding clinical efficacy for COVID-19 are being evaluated.(138)
 - Other fibrinolytics for which COVID-19 efficacy data are being evaluated include defibrotide.(139) (140)
- Safety Concerns: (136) (137)
 - Bleeding

Inhaled Pulmonary Vasodilators:

- There is no evidence for routine use of inhaled pulmonary vasodilators (e.g., nitric oxide, prostacyclins) in acute respiratory failure in COVID-19 patients. Avoid aerosolized vasodilators.(26) (60) (61) (133)
- Use may be considered in specific patients with ARDS as a temporizing measure when patients develop refractory hypoxemia despite optimization of ventilation and other rescue strategies.(26) (60)
- If inhaled pulmonary vasodilator therapy is used, a short trial with preestablished criteria for ongoing use or discontinuation is recommended.(26) (61) (133)
- Additional data regarding clinical efficacy for COVID-19 are being evaluated.(58) (59)

Interleukin-1 (IL-1) Antagonists:

- Rationale for Use: Cytokine release syndrome may be a component of severe disease in COVID-19 patients.(24) (89) (90) (107)
- Mechanism of Action: Interleukin-1 antagonists, such as anakinra and canakinumab, prevent the binding of IL-1 (a pro-inflammatory cytokine that mediates various inflammatory and immunological responses, including activation of IL-6) to interleukin-1 receptors. Anakinra acts similarly to the native interleukin-1 receptor antagonist by competitively inhibiting the binding of both IL-1 alpha and IL-1 beta to the IL-1 type 1 receptor. Canakinumab is a human monoclonal antibody that specifically targets and neutralizes IL-1 beta; thereby preventing its interaction with IL-1 receptors.(89) (106) (147)
- Evidence / Experience:
 - Due to a lack of clinical data, the NIH COVID-19 treatment guidelines do not give recommendations for or against the use of IL-1 antagonists.(133)
 - Anakinra:
 - Retrospective cohort study comparing anakinra plus standard therapy to standard therapy alone in patients with COVID-19, moderate-to-severe ARDS, and hyperinflammation.(148)
 - 21-day survival was 90% in the anakinra group and 56% in standard treatment group (p = 0.009). Respiratory function improved in 72% (n = 21/29) of anakinra patients and 50% (n = 8/16) of patients in the standard treatment group.
 - Additional data regarding clinical efficacy for COVID-19 are being evaluated.(132)
 - Other IL-1 antagonists for which COVID-19 efficacy data are being evaluated include canakinumab.(132)
- Safety Concerns: (106) (147)
 - Caution in patients with thrombocytopenia and neutropenia
 - Infusion-related reactions (anakinra)

Interleukin-6 (IL-6) Receptor Antagonists:

- Rationale for Use: Cytokine release syndrome may be a component of severe disease in COVID-19 patients.(24) (25) (89) (90)
- Mechanism of Action: IL-6 receptor-inhibiting monoclonal antibodies block IL-6-mediated signaling by competitively binding to both soluble and membrane-bound IL-6 receptors (sIL-6R and mIL-6R). IL-6 is a proinflammatory cytokine that is involved in diverse physiological processes such as T-cell activation, immunoglobulin secretion induction, hepatic acute-phase protein synthesis initiation, and hematopoietic precursor cell proliferation and differentiation stimulation. IL-6 is produced by various cell types, including T- and B-cells, lymphocytes, monocytes, and fibroblasts.(52) (72) (78) (97) (98)
- Evidence / Experience:
 - Due to a lack of clinical data, the NIH COVID-19 treatment guidelines do not recommend for or against the use of IL-6 receptor inhibitors.(133)
 - Siltuximab:
 - A retrospective study of 21 patients with COVID-19 induced pneumonia/ARDS analyzed patients who received treatment with siltuximab.(101)
 - CRP concentrations reduced to within normal range by day 5 and remained stable in all 16 patients with available data; 33% (n = 7/21) condition improved with reduced need for ventilation; 43% (n = 9/21) condition stabilized; 24% (n = 5/21) condition worsened and required intubation.
 - A cohort study of patients treated with standard therapy is ongoing
 - Additional data regarding clinical efficacy for COVID-19 are being evaluated.(132)
 - Tocilizumab:
 - A retrospective review analyzed 21 patients in which tocilizumab was added to standard COVID-19 therapy.(25)
 - Preliminary data suggest tocilizumab may have clinical benefit as adjunctive therapy.
 - Clinical symptoms, CT opacity changes, lymphocyte percentage, and CRP concentrations all improved in these patients; however, no comparators were reported.
 - Some protocols include recommendations for use.(21)
 - Additional data regarding clinical efficacy for COVID-19 are being evaluated.(132)
 - Other IL-6 receptor inhibitors for which COVID-19 efficacy data are being evaluated include clazakizumab and sarilumab.(132)
- Safety Concerns: (52) (78) (97)
 - Risk of GI perforation
 - Risk of hepatotoxicity
 - Caution in patients with thrombocytopenia and neutropenia
 - Infusion-related reactions

Janus Kinase (JAK) Inhibitors:

- Rationale for Use: Cytokine release syndrome may be a component of severe disease in COVID-19 patients.(24) (89) (90) (94)
- Mechanism of Action: Janus kinases are intracellular enzymes that transmit signals arising from the interaction of cytokines and growth factors with receptors located on the cellular membrane. These enzymes phosphorylate and activate signal transducers and activators of transcription proteins (STATs), which modulate intracellular activity including gene expression. The JAK-mediated signaling pathway is pivotal in influencing immune system activation, as cytokine receptors are expressed on most immune cells. JAK inhibitors modulate the signaling pathway by preventing the phosphorylation and activation of STATs.(92) (130) (131)
- Evidence / Experience:
 - Due to the broad immunosuppressive effect, the NIH COVID-19 treatment guidelines recommend against the use of JAK inhibitors outside of clinical trials.(133)
 - Baricitinib:
 - A non-randomized, open-label trial compared safety and efficacy of baricitinib plus lopinavir; ritonavir in 12 patients with moderate COVID-19 against a control group treated with hydroxychloroquine plus lopinavir; ritonavir.(144)
 - For baricitinib-treated patients, all clinical characteristics and respiratory function parameters [i.e., fever, SpO₂, PaO₂/FiO₂, CRP, Modified Early Warning Score (MEWS)] improved at weeks 1 and 2 compared to baseline. For control group, no significant changes were noted at week 2 compared to baseline.
 - Preliminary results found no serious adverse events reported in the baricitinib-treated patients; however, 1 patient stopped treatment after 10 days due to increased liver function test.
 - Additional data regarding clinical efficacy for COVID-19 are being evaluated.(132)
 - Other JAK inhibitors for which COVID-19 efficacy data are being evaluated include: ruxolitinib and tofacitinib.(132)
- Safety Concerns: (92) (130) (131)
 - Thrombosis, including deep vein thrombosis (DVT) and pulmonary embolism (PE)
 - Risk of GI perforation
 - Caution in patients with neutropenia, lymphopenia, and anemia
 - Monitor for elevated liver function tests (LFTs)

NSAIDs:

- The NIH COVID-19 treatment guidelines recommend there be no difference in the use of antipyretic treatments (e.g., acetaminophen or NSAIDs) between patients with or without COVID-19. Patients taking NSAIDs for comorbid conditions should continue therapy as previously directed by their prescriber.(133)
- ESICM and SCCM Surviving Sepsis Campaign recommendations suggest acetaminophen for temperature control in critically ill adults with COVID-19 who develop fever.(26)
- The FDA continues to investigate the use of NSAIDs in patients with COVID-19 symptoms.(20)

- Concern for potential worsening of COVID-19 symptoms has been suggested, but confirmatory clinical data is lacking at this time.(5)
- There is an anecdotal published letter that suggests a link between ibuprofen and increased ACE2 expression that may lead to worse outcomes in COVID-19 patients.(50)

Nutritional Supplements

- The role of nutritional supplements for the treatment or prevention of COVID-19 is unknown. Several supplements are under investigation in combination with other treatment modalities (e.g. zinc, vitamin C, vitamin D) for both treatment and prophylaxis.(111) (112) (113) (114) (115) (116) (117) (118) (119) (120)
- Safety concerns include adverse events from large doses and the potential for drug interactions.(108) (109) (110)

Understanding of the treatment of patients with COVID-19 is rapidly evolving. Information will continue to emerge regarding pharmacologic therapy for SARS-CoV-2 as clinical data are reported.

References:

1. World Health Organization. Clinical management of severe acute respiratory infection when novel coronavirus (nCoV) infection is suspected [https://www.who.int/publications-detail/clinical-management-of-severe-acute-respiratory-infection-when-novel-coronavirus-\(ncov\)-infection-is-suspected](https://www.who.int/publications-detail/clinical-management-of-severe-acute-respiratory-infection-when-novel-coronavirus-(ncov)-infection-is-suspected).
2. CDC Website: <https://www.cdc.gov/coronavirus/2019-ncov/hcp/clinical-guidance-management-patients.html>.
3. FDA Website: <https://www.fda.gov/emergency-preparedness-and-response/mcm-issues/coronavirus-disease-2019-covid-19>.
4. Chu CM, Cheng VCC, Hung IFN, et al. Role of lopinavir/ritonavir in the treatment of SARS: Initial virological and clinical findings. *Thorax* 2004;59(3):252–256. PMID: 1498565
5. Loutfy MR, Blatt LM, Siminovitch KA, et al. Interferon Alfacon-1 Plus Corticosteroids in Severe Acute Respiratory Syndrome: A Preliminary Study. *J Am Med Assoc* 2003;290(24):3222–3228. PMID: 14693875
6. Peiris JSM, Chu CM, Cheng VCC, et al. Clinical progression and viral load in a community outbreak of coronavirus-associated SARS pneumonia: A prospective study. *Lancet* 2003;361(9371):1767–1772. PMID: 12781535
7. Jin Y., Cai, L., Cheng, Z. et al. A rapid advice guideline for the diagnosis and treatment of 2019 novel coronavirus (2019-nCoV) infected pneumonia (standard version). *Military Med Res* 7, 4 (2020).
8. Huang C, Wang Y, Li X, et al. Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet* Published online January 24, 2020. PMID: 31986264
9. ClinicalTrials.gov. National Institutes of Health (NIH) U.S. National Library of Medicine
ClinicalTrials.gov website
https://www.clinicaltrials.gov/ct2/results?cond=Coronavirus&term=&type=&rslt=&age_v=&gndr=&intr

[=remdesivir&titles=&outc=&spons=&lead=&id=&cntry=&state=&city=&dist=&locn=&rsub=&strd_s=&trd_e=&prcd_s=&prcd_e=&sfpd_s=&sfpd_e=&rfpd_s=&rfpd_e=&lupd_s=&lupd_e=&sort=](#)

10. Agostini ML, Andres EL, Sims AC, et al. Coronavirus susceptibility to the antiviral remdesivir (GS-5734) is mediated by the viral polymerase and the proofreading exoribonuclease. *MBio* 2018;9(2):1–15. PMID: 29511076
11. Elfiky AA. Anti-HCV, nucleotide inhibitors, repurposing against COVID-19. *Life Sciences* 2020 May 1;248:117477. PMID: 32119961
12. Korea Biomedical Review website: <http://www.koreabiomed.com/news/articleView.html?idxno=7428>.
13. Gao J, T Zhenxue, Yang X. Breakthrough: Chloroquine phosphate has shown apparent efficacy in treatment of COVID-19 associated pneumonia in clinical studies. *Biosci Trends* 2020;14(1):72-73. PMID: 32074550
14. Wang M, Cao R, Zhang L, et al. Remdesivir and chloroquine effectively inhibit the recently emerged novel coronavirus (2019-nCoV) in vitro. *Cell Research* 2020;30:269–271. PMID: 32020029
15. Yao X, Ye F, Zhang M, et al. In vitro antiviral activity and projection of optimized dosing design of hydroxychloroquine for the treatment of Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2). *Clin Infect Dis* 2020 Mar 9. [Epub ahead of print] PMID: 32150618
16. World Health Organization (WHO). Coronavirus: landscape analysis of therapeutics as of 17 February 2020. Retrieved March 16, 2020. Available on the World Wide Web at https://www.who.int/blueprint/priority-diseases/key-action/Table_of_therapeutics_Appendix_17022020.pdf?ua=1.
17. Colson P, Rolain J, Lagier J, et al. Chloroquine and hydroxychloroquine as available weapons to fight COVID-19. *Int J Antimicrob Agents* 2020. [Epub ahead of print] PMID: 32145363
18. Dong L, Hu S, Gao J. Discovering drugs to treat coronavirus disease 2019 (COVID-19). *Drug Discov Ther* 2020;14:58-60. PMID: 32147628
19. Multicenter Collaboration Group of Department of Science and Technology of Guangdong Province and Health Commission of Guangdong Province for chloroquine in the treatment of novel coronavirus pneumonia. Expert consensus on chloroquine phosphate for the treatment of novel coronavirus pneumonia. *Zhonghua Jie He Hu Xi Za Zhi* 2020 Mar;43:185-188. PMID: 32164085
20. FDA Website: <https://www.fda.gov/drugs/drug-safety-and-availability/fda-advises-patients-use-non-steroidal-anti-inflammatory-drugs-nsaids-covid-19>.
21. Italian Society of Infectious and Tropical Diseases. Handbook for the care of people with disease-COVI 19. Edition 2.0, March 13, 2020.
22. FDA Website: https://www.fda.gov/vaccines-blood-biologics/investigational-new-drug-ind-or-device-exemption-ide-process-cber/investigational-covid-19-convalescent-plasma-emergency-inds?utm_campaign=What%27sNew2020-03-24&utm_medium=email&utm_source=Eloqua.
23. Cao B, Wang Y, Wen D, et al. A trial of lopinavir-ritonavir in adults hospitalized with severe Covid-19. *N Engl J Med* 2020;382(19):1787-1799. PMID: 32187464
24. Zhou F, Yu T, Du R, et al. Clinical course and risk factors for mortality of adult inpatients with COVID-19 in Wuhan, China: a retrospective cohort study. *Lancet* 2020 Mar 11. [Epub ahead of print] PMID: 32171076.
25. Xu X, Han M, LI T, et al. Effect treatment of severe COVID-19 patients with tocilizumab. *ChinaXiv.20200300026.v1*
26. ESICM, SCCM. Surviving sepsis campaign rapid guidelines of the management of critically ill adults with coronavirus disease 2019 (pre-publication). Available on the World Wide Web at: <https://www.esicm.org/ssc-covid19-guidelines/>.

27. Gautret P, Lagier J, Parola P, et al. Hydroxychloroquine and azithromycin as a treatment of COVID-19: results of an open-label non-randomized clinical trial. *Int J Antimicrob Agents* 2020 Mar 20. [Epub ahead of print] PMID: 32205204
28. Arabi YM, Deeb A, Al-Hameed, et al. Macrolides in critically ill patients with Middle East Respiratory Syndrome. *Int J Infect Dis* 2019;81:184-190. PMID: 30690213
29. Fox R. Anti-malarial drugs: possible mechanisms of action in autoimmune disease and prospects for drug development. *Lupus* 1996;5 Suppl 1:S4-10. PMID: 8803903
30. Ben-Zvi H, Kivity S, Langevitz P, et al. Hydroxychloroquine: from malaria to autoimmunity. *Clin Rev Allergy Immunol* 2012;42:145-153. PMID: 21221847
31. Cortegiani A, Ingoglia G, Ippolito M, et al. A systematic review on the efficacy and safety of chloroquine for the treatment of COVID-19. *J Crit Care* 2020 Mar 10. [Epub ahead of print] PMID: 32173110
32. Savarino A, Trani LD, Donatelli I, et al. New insights into the antiviral effects of chloroquine. *The Lancet* 2006;6:67-9. PMID: 16439323
33. Fox RI. Mechanism of action of hydroxychloroquine as an antirheumatic drug. *Semin Arthritis Rheum* 1993; 23(2 Suppl 1):82-91. PMID: 8278823
34. Amsden GW. Anti-inflammatory effects of macrolides - an underappreciated benefit in the treatment of community-acquired respiratory tract infections and chronic inflammatory pulmonary conditions? *J Antimicrob Chemother* 2005;55:10-21. PMID: 15590715
35. Beigelman A, Mikols CL, Gunsten SP, et al. Azithromycin attenuates airway inflammation in a mouse model of viral bronchiolitis. *Respir Res* 2010;11:90. PMID: 20591166
36. Kanoh S, Rubin BK. Mechanisms of action and clinical application of macrolides as immunomodulatory medications. *Clin Microbiol Rev* 2010;23:590-615. PMID: 20610825
37. Zarogoulidis P, Papanas N, Kioumis I, et al. Macrolides: from in vitro anti-inflammatory and immunomodulatory properties to clinical practice in respiratory disease. *Eur J Clin Pharmacol* 2012;68:479-503. PMID: 22105373
38. U.S. Army Medical Research and Development Command. Expanded access remdesivir (RDV; GS-5734). Retrieved March 18, 2020. Available on the World Wide Web at: <https://clinicaltrials.gov/ct2/show/NCT04302766?term=remdesivir&draw=2&rank=3>.
39. Brown AJ, Won JJ, Graham RL, et al. Broad spectrum antiviral remdesivir inhibits human endemic and zoonotic deltacoronaviruses with a highly divergent RNA dependent RNA polymerase. *Antiviral Research* 2019;169:1-10. PMID: 31233808
40. Regeneron Pharmaceuticals. Evaluation of the efficacy and safety of sarilumab in hospitalized patients with COVID-19. Retrieved March 24, 2020. Available on the World Wide Web at: <https://clinicaltrials.gov/ct2/show/NCT04315298?term=sarilumab&draw=3&rank=4>.
41. de Wit, Feldmann F, Cronin J, et al. Prophylactic and therapeutic remdesivir (GS-5734) treatment in the rhesus macaque model of MERS-CoV infection. *Proc Natl Acad Sci U S A*. 2020 Feb 13.[Epub ahead of print] PMID: 32054787
42. Ko W, Rolain J, Lee N, et al. Arguments in favor of remdesivir for treating SARS-CoV-2 infections. *Int J Antimicrob Agents* 2020 Mar 6. [Epub ahead of print] PMID: 32147516
43. Gordon CJ, Tchesnokov EP, Feng JY, et al. The antiviral compound remdesivir potently inhibits RNA-dependent RNA polymerase from Middle East respiratory syndrome coronavirus. *J Biol Chem* 2020 Feb 24. [Epub ahead of print] PMID: 32094225
44. Warren TK, Jordan R, Lo MK, et al. Therapeutic efficacy of the small molecule GS-5734 against Ebola virus in rhesus monkeys. *Nature* 2016;531:381–385. PMID: 26934220
45. Kaletra (lopinavir; ritonavir) tablet and solution package insert. North Chicago, IL: AbbVie Inc; 2019 Dec.

46. Aralen (chloroquine) package insert. Bridgewater, NJ: Sanofi-aventis U.S. LLC.; 2018 Oct.
47. Plaquenil (hydroxychloroquine) package insert. St. Michael, Barbados: Concordia Pharmaceuticals, Inc.; 2017 Jan.
48. Zithromax (azithromycin 250 mg and 500 mg tablets and azithromycin oral suspension) package insert. New York, NY: Pfizer Inc.; 2019 Apr.
49. Credible Meds. COVID-19 experimental therapies and TdP risk. Retrieved March 24, 2020. Available on the World Wide Web at: <https://crediblemeds.org/blog/covid-19-experimental-therapies-and-tdp-risk/>
50. Fang L, Karakiulakis G, Roth M. Are patients with hypertension and diabetes mellitus at increased risk for COVID-19 infection? *Lancet Respir Med* 2020 Mar 11. [Epub ahead of print] PMID:32171062
51. Peking University First Hospital. Favipiravir combined with tocilizumab in the treatment of Corona Virus Disease 2019. Retrieved March 24, 2020. Available on the World Wide Web at: <https://clinicaltrials.gov/ct2/show/NCT04310228?cond=Coronavirus&intr=Tocilizumab&draw=2&rank=1>.
52. Actemra (tocilizumab) injection package insert. South San Francisco, CA: Genentech, Inc.; 2019 Jun.
53. Chen F, Chan KH, Jiang Y et al. In vitro susceptibility of 10 clinical isolates of SARS coronavirus to selected antiviral compounds. *J Clin Virol* 2004; 31:69-75. PMID: 15288617
54. Yao TT, Qian JD, Zhu WY et al. A systematic review of lopinavir therapy for SARS coronavirus and MERS coronavirus-A possible reference for coronavirus disease-19 treatment option. *J Med Virol* 2020 Feb 27. [Epub ahead of print] PMID: 32104907
55. Liu X, Wang XJ. Potential inhibitors for 2019-nCoV coronavirus M protease from clinically proven medicines. *J Genet Genomics* 2020 Feb 13. [Epub ahead of print] PMID: 32173287
56. Chen J, Liu D, Liu L, et al. A pilot study of hydroxychloroquine in treatment of patients with common coronavirus disease-19 (COVID-19). *J Zhejiang Univ (Med Sci)* 2020:49(1).
57. Institute for Safe Medication Practices (ISMP). Acute Care ISMP Medication Safety Alert. Special edition COVID-19. 2020 March;25(6):1-5.
58. Massachusetts General Hospital. Nitric oxide gas inhalation therapy for mild/moderate COVID-19 (NoCovid). Retrieved March 30, 2020. Available on the World Wide Web at: <https://clinicaltrials.gov/ct2/show/NCT04305457?cond=Coronavirus&intr=Nitric+Oxide&draw=2&rank=4>.
59. Massachusetts General Hospital. Nitric oxide gas inhalation for Severe Acute Respiratory Syndrome in COVID-19 (NOSARSCOVID). Retrieved March 30, 2020. Available on the World Wide Web at: <https://clinicaltrials.gov/ct2/show/NCT04306393?cond=Coronavirus&intr=Nitric+Oxide&draw=2&rank=2>.
60. Department of Defense (DoD). DoD COVID-19 practice management guide: clinical management of COVID-19. March 23, 2020. Available on the World Wide Web at: <https://www.health.mil/Reference-Center/Technical-Documents/2020/03/24/DoD-COVID-19-Practice-Management-Guide>
61. American Association for Respiratory Care (AARC). SARS CoV-2 guidance document. Retrieved March 30, 2020. Available on the World Wide Web at: <https://www.aarc.org/wp-content/uploads/2020/03/guidance-document-SARS-COVID19.pdf>.
62. Wu C, Chen X, Cai Y, et al. Risk factors associated with acute respiratory distress syndrome and death in patients with coronavirus disease 2019 pneumonia in Wuhan, China. *JAMA Intern Med*. 2020 Mar 13. [Epub ahead of print] PMID: 32167524
63. Zhou YH, Qin YY, Lu YQ, et al. Effectiveness of glucocorticoid therapy in patients with severe novel coronavirus pneumonia: protocol of a randomized controlled trial. *Chin Med J (Engl)*. 2020 Mar 5. [Epub ahead of print] PMID: 32149773

64. Lian J, Jin X, Hao S, et al. Analysis of epidemiological and clinical features in older patients with Corona Virus Disease 2019 (COVID-19) out of Wuhan. Clin Infect Dis. 2020 Mar 25. [Epub ahead of print] PMID: 32211844
65. Shen C, Wang Z, Zhao F, et al. Treatment of 5 critically ill patients with COVID-19 with convalescent plasma. JAMA 2020 Mar 27. [Epub ahead of print] PMID: 32219428
66. Food and Drug Administration (FDA). Chloroquine phosphate or hydroxychloroquine sulfate supplied from the strategic national stockpile for treatment of 2019 coronavirus disease: emergency use authorization letter. Retrieved March 30, 2020. Available on the World Wide Web at: <https://www.fda.gov/emergency-preparedness-and-response/mcm-legal-regulatory-and-policy-framework/emergency-use-authorization#2019-ncov>.
67. Food and Drug Administration (FDA). Fact sheet for health care providers: emergency use authorization (EUA) of chloroquine phosphate supplied from the strategic national stockpile for treatment of COVID-19 in certain hospitalized patients. Retrieved March 30, 2020. Available on the World Wide Web at: <https://www.fda.gov/emergency-preparedness-and-response/mcm-legal-regulatory-and-policy-framework/emergency-use-authorization#2019-ncov>.
68. Food and Drug Administration (FDA). Fact sheet for health care providers: emergency use authorization (EUA) of hydroxychloroquine sulfate supplied from the strategic national stockpile for treatment of COVID-19 in certain hospitalized patients. Retrieved March 30, 2020. Available on the World Wide Web at: <https://www.fda.gov/emergency-preparedness-and-response/mcm-legal-regulatory-and-policy-framework/emergency-use-authorization#2019-ncov>.
69. Chen Z, Hu J, Zhang Z, et al. Efficacy of hydroxychloroquine in patients with COVID-19: results of a randomized clinical trial. 2020 DOI: <https://doi.org/10.1101/2020.03.22.20040758>
70. Miao M, De Clercq E, Li G. Clinical significance of chemokine receptor antagonists. Expert Opin Drug Metab Toxicol. 2020 Jan;16(1):11-30. [Epub ahead of print] PMID: 31903790
71. CytoDyn. Press release. Retrieved April 2, 2020. Available on the World Wide Web at: <https://www.cytodyn.com/newsroom/press-releases/detail/405/treatment-with-cytodyns-leronlimab-indicates-significant>
72. Assistance Publique – Hopitaux de Paris. Cohort multiple randomized controlled trials open-label of immune modulatory drugs and other treatments in COVID-19 patients – sarilumab trial – CORIMUNO-19-SARI (CORIMUNO-SARI). Retrieved April 2, 2020. Available on the World Wide Web at: <https://clinicaltrials.gov/ct2/show/NCT04324073?term=sarilumab&draw=3&rank=2>
73. Peking University First Hospital. Favipiravir combined with tocilizumab in the treatment of Corona Virus Disease 2019. Retrieved March 24, 2020. Available on the World Wide Web at: <https://clinicaltrials.gov/ct2/show/NCT04310228?cond=Coronavirus&intr=Tocilizumab&draw=2&rank=1>
74. Beijing Chao Yang Hospital. Clinical trial of favipiravir tablets combine with chloroquine phosphate in treatment of novel coronavirus pneumonia. Retrieved April 2, 2020. Available on the World Wide Web at: <https://clinicaltrials.gov/ct2/show/NCT04319900?term=favipiravir&draw=2&rank=9>
75. Shiraki K and Daikoku T. Favipiravir, an anti-influenza drug against life-threatening RNA virus infections. Pharmacol Ther. 2020 Feb 22. [Epub ahead of print] PMID: 32097670
76. Regneron. Press release. Retrieved April 2, 2020. Available on the World Wide Web at: <https://newsroom.regeneron.com/news-releases/news-release-details/regeneron-and-sanofi-begin-global-kevzara-sarilumab-clinical>
77. Sanofi. Press release. Retrieved April 2, 2020. Available on the World Wide Web at: <http://www.news.sanofi.us/2020-03-16-Sanofi-and-Regeneron-begin-global-Kevzara-R-sarilumab-clinical-trial-program-in-patients-with-severe-COVID-19>
78. Kevzara (sarilumab) package insert. Bridgewater, NJ: Sanofi-Aventis US. LLC; 2018 Apr.

79. Institute for Safe Medication Practices (ISMP). Acute Care ISMP Medication Safety Alert. 2020 April;25(6 Supplement):1-5.
80. Thachil J, Tang N, Gando S, et al. ISTH interim guidance on recognition and management of coagulopathy in COVID-19. *J Thromb Haemost* 2020 [Epub ahead of print] DOI: doi:10.1111/jth.14810.
81. Tang N, Bai H, Chen X. Anticoagulant treatment is associated with decreased mortality in severe coronavirus disease 2019 patients with coagulopathy. *J Thromb Haemost*. 2020. [Epub ahead of print] PMID: 32220112
82. Regeneron Pharmaceuticals. Evaluation of the efficacy and safety of sarilumab in hospitalized patients with COVID-19. Retrieved April 2, 2020: Available on the World Wide Web at: <https://clinicaltrials.gov/ct2/show/study/NCT04315298?term=sarilumab&draw=2>
83. Sanofi. Sarilumab COVID-19. Retrieved April 2, 2020. Available on the World Wide Web at: <https://clinicaltrials.gov/ct2/show/NCT04327388?cond=Coronavirus&intr=sarilumab&draw=2&rank=2>
84. Henriksen M. Anti-il6 treatment of serious COVID-19 disease with threatening respiratory failure (TOCIDVID). Retrieved April 2, 2020. Available on the World Wide Web at: <https://clinicaltrials.gov/ct2/show/NCT04322773?cond=Coronavirus&intr=sarilumab&draw=2&rank=3>
85. Barrett L. Treatment of moderate to severe coronavirus disease (COVID-19) in hospitalized patients. Retrieved April 2, 2020. Available on the World Wide Web at: <https://clinicaltrials.gov/ct2/show/NCT04321993?cond=Coronavirus&intr=sarilumab&draw=2&rank=4>
86. Assistance Publique - Hopitaux de Paris. Cohort multiple randomized controlled trials open-label of immune modulatory drugs and other treatments in COVID-19 patients - tocilizumab trial - CORIMUNO-19 - TOCI (CORIMUNO-TOCI) (CORIMUNO-TOC). Retrieved April 2, 2020. Available on the World Wide Web at: <https://clinicaltrials.gov/ct2/show/NCT04331808?cond=Coronavirus&intr=Tocilizumab&draw=2&rank=1>
87. University Hospital Ghent. Treatment of COVID-19 patients with anti-interleukin drugs (COV-AID). Retrieved April 2, 2020. Available on the World Wide Web at: <https://clinicaltrials.gov/ct2/show/NCT04330638?cond=Coronavirus&intr=Tocilizumab&draw=2&rank=4>
88. Molina JM, Delaugerre C, Goff JL, et al. No evidence of rapid antiviral clearance or clinical benefit with the combination of hydroxychloroquine and azithromycin in patients with severe COVID-19. *Med Mal Infect* 2020. [Epub ahead of print] DOI:https://doi.org/10.1016/j.medmal.2020.03.006. PMID: 32240719
89. Zhang C, Wu Z, Li JW, et al. The cytokine release syndrome (CRS) of severe COVID-19 and Interleukin-6 receptor (IL-6R) antagonist tocilizumab may be the key to reduce the mortality. *Int J Antimicrob Agents*. 2020. [Epub ahead of print] PMID: 32234467
90. Zhang W, Zhao Y, Zhang F, et al. The use of anti-inflammatory drugs in the treatment of people with severe coronavirus disease 2019 (COVID-19): the perspectives of clinical immunologists from China. *Clin Immunol*. 2020. [Epub ahead of print] PMID: 32222466
91. Lin L, Lu L, Cao W, et al. Hypothesis for potential pathogenesis of SARS-CoV-2 infection—a review of immune changes in patients with viral pneumonia. *Emerg Microbes Infect*. 2020;9:727-732. PMID: 32196410
92. Olumiant (baricitinib) tablets package insert. Indianapolis, IN: Lilly USA, LLC; 2019 Oct.
93. Barret L. Treatment of moderate to severe coronavirus disease (COVID-19) in hospitalized patients. Retrieved April 2, 2020. Available on the World Wide Web at: <https://clinicaltrials.gov/ct2/show/NCT04321993?cond=Coronavirus&intr=Baricitinib&draw=2&rank=1>

94. Stebbing J, Phelan A, Griffin I, et al. COVID-19: combining antiviral and anti-inflammatory treatments. *Lancet Infect Dis*. 2020. [Epub ahead of print] PMID: 32113509
95. American Society of Hematology (ASH). COVID-19 and VTE/anticoagulation: frequently asked questions Version 2.1. Retrieved April 29, 2020. Available on the World Wide Web at: <https://www.hematology.org/covid-19/covid-19-and-vte-anticoagulation>
96. Hospital of Prato. Baricitinib in symptomatic patients infected by COVID-19: an open-label, pilot study (BARI-COVID). Retrieved April 6, 2020. Available on the World Wide Web at: <https://clinicaltrials.gov/ct2/show/NCT04320277?cond=COVID&intr=Baricitinib&draw=2&rank=1>
97. Sylvant (siltuximab) package insert. Hemel Hempstead, United Kingdom: EUSA Pharma. Ltd; 2019 Dec.
98. Maston LD, Jones DT, Giermakowska W, et al. Interleukin-6 trans-signaling contributes to chronic hypoxia-induced pulmonary hypertension. *Pulm Circ*. 2018;8(3):1-11. PMID: 29767573
99. Martinez JP. Efficacy and safety of siltuximab vs. corticosteroids in hospitalized patients with COVID19 pneumonia. Retrieved April 8, 2020. Available on the World Wide Web at: <https://clinicaltrials.gov/ct2/show/NCT04329650?term=Siltuximab&draw=3&rank=12>
100. A.O. Ospedale Papa Giovanni XXIII. An observational case-control study of the use of siltuximab in ARDS patients diagnosed with COVID-19 infection (SISCO). Retrieved April 8, 2020. Available on the World Wide Web at: <https://clinicaltrials.gov/ct2/show/NCT04322188?term=Siltuximab&draw=3&rank=13>
101. Gritti G, Raimondi F, Ripamonti D, et al. Use of siltuximab in patients with COVID-19 pneumonia requiring ventilatory support. Retrieved April 8, 2020. Available on the World Wide Web at: <https://www.medrxiv.org/content/10.1101/2020.04.01.20048561v1>
102. American Society of Hematology (ASH). COVID-19 and coagulopathy: frequently asked questions Version 2.0. Retrieved April 22, 2020. Available on the World Wide Web at: <https://www.hematology.org/covid-19/covid-19-and-coagulopathy>
103. Hunt B, Retter A, McClintock C. Practical guidance for the prevention of thrombosis and management of coagulopathy and disseminated intravascular coagulation of patients infected with COVID-19. Retrieved April 9, 2020. Available on the World Wide Web at: <https://thrombosisuk.org/covid-19-thrombosis.php>
104. Swedish Orphan Biovitrum. Efficacy and safety of emapalumab and anakinra in reducing hyperinflammation and respiratory distress in patients with COVID-19 infection. Retrieved April 9, 2020. Available on the World Wide Web at: <https://clinicaltrials.gov/ct2/show/NCT04324021>
105. Hellenic Institute for the study of sepsis. Personalized immunotherapy for SARS-CoV-2 (COVID-19) associated with organ dysfunction (ESCAPE). Retrieved April 9, 2020. Available on the World Wide Web: <https://clinicaltrials.gov/ct2/show/NCT04339712?term=anakinra&cond=covid&draw=2&rank=2>
106. Kineret (anakinra) package insert. Stockholm, Sweden: Swedish Orphan Biovitrum AB; 2018 Jun.
107. Mehta P, McAuley DF, Brown M, et al. COVID-19: consider cytokine storm syndromes and immunosuppression. *Lancet*. 2020;395(10229):1033-1034. PMID: 32192578
108. National Institutes of Health (NIH). Zinc: fact sheet for health professionals. Retrieved April 9, 2020. Available on the World Wide Web at: <https://ods.od.nih.gov/factsheets/Zinc-HealthProfessional/>
109. National Institutes of Health (NIH). Vitamin C: fact sheet for health professionals. Retrieved April 9, 2020. Available on the World Wide Web at: <https://ods.od.nih.gov/factsheets/VitaminC-HealthProfessional/>
110. National Institutes of Health (NIH). Vitamin D: fact sheet for health professionals. Retrieved April 9, 2020. Available on the World Wide Web at: <https://ods.od.nih.gov/factsheets/VitaminD-HealthProfessional/>

111. ProgenaBiome. A study of hydroxychloroquine, vitamin C, vitamin D, and zinc for the prevention of COVID-19 infection (HELPCOVID-19). Retrieved April 9, 2020. Available on the World Wide Web at: <https://clinicaltrials.gov/ct2/show/NCT04335084?cond=COVID&intr=zinc&draw=2&rank=2>
112. Istinye University. Proflaxis using hydroxychloroquine plus vitamins-zinc during COVID-19 pandemia. Retrieved April 9, 2020. Available on the World Wide Web at: <https://clinicaltrials.gov/ct2/show/NCT04326725?cond=COVID&intr=zinc&draw=2&rank=3>
113. ProgenaBione. A study of quintuple therapy to treat COVID-19 infection (HAZCpaC). Retrieved April 9, 2020. Available on the World Wide Web at: <https://clinicaltrials.gov/ct2/show/NCT04334512?cond=COVID&intr=zinc&draw=2&rank=4>
114. University of Melbourne. World-first trial to test benefit of intravenous zinc in COVID-19 fight. Retrieved April 9, 2020. Available on the World Wide Web at: <https://medicalxpress.com/news/2020-04-world-first-trial-benefit-intravenous-zinc.html>
115. University of Palermo. Use of ascorbic acid in patients With COVID 19. Retrieved April 9, 2020. Available on the World Wide Web at: <https://clinicaltrials.gov/ct2/show/NCT04323514?cond=COVID&intr=vitamin+C&draw=2&rank=1>
116. Providence Health and Services. Hydroxychloroquine in patients with newly diagnosed COVID-19 compared to standard of care. Retrieved April 9, 2020. Available on the World Wide Web at: <https://clinicaltrials.gov/ct2/show/NCT04334967?cond=COVID&intr=vitamin+C&draw=2&rank=4>
117. Washington University. Hydroxychloroquine for COVID-19 PEP. Retrieved April 9, 2020. Available on the World Wide Web at: <https://clinicaltrials.gov/ct2/show/NCT04328961?cond=COVID&intr=vitamin+C&draw=2&rank=5>
118. ZhiYong Peng. Vitamin C infusion for the treatment of severe 2019-nCoV infected pneumonia. Retrieved April 9, 2020. Available on the World Wide Web at: <https://clinicaltrials.gov/ct2/show/NCT04264533?cond=COVID&intr=vitamin+C&draw=2&rank=6>
119. Universite de Sherbrooke. Lessening organ dysfunction with vitamin C (LOVIT). Retrieved April 9, 2020. Available on the World Wide Web at: <https://clinicaltrials.gov/ct2/show/NCT03680274?cond=COVID&intr=vitamin+C&draw=2&rank=8>
120. Universidad de Granada. Vitamin D on Prevention and Treatment of COVID-19 (COVITD-19). Retrieved April 9, 2020. Available on the World Wide Web at: <https://clinicaltrials.gov/ct2/show/NCT04334005?cond=COVID&intr=Vitamin+D&draw=2&rank=1>
121. University of Colorado. Safety and efficacy of baricitinib for COVID-19. Retrieved April 14, 2020. Available on the World Wide Web at: <https://clinicaltrials.gov/ct2/show/NCT04340232?cond=COVID&intr=Baricitinib&draw=2&rank=1>
122. Benfield T. Efficacy and safety of novel treatment options for adults with COVID-19 pneumonia (CCAP). Retrieved April 14, 2020. Available on the World Wide Web at: <https://clinicaltrials.gov/ct2/show/NCT04345289?cond=COVID&intr=Baricitinib&draw=2&rank=4>
123. Assistance Publique – Hopitaux de Paris. Study of Immune Modulatory Drugs and Other Treatments in COVID-19 Patients: Sarilumab, Azithromycin, Hydroxychloroquine Trial - CORIMUNO-19 - VIRO (CORIMUNO-VIRO). Retrieved April 13, 2020. Available on the World Wide Web at: <https://clinicaltrials.gov/ct2/show/NCT04341870?term=sarilumab&cond=SARS-CoV+2&draw=2&rank=1>
124. Grein J, Ohmagari N, Shin D, et al. Compassionate use of remdesivir for patients with severe Covid-19. *New Engl J Med* 2020. [Epub ahead of print] PMID: 32275812
125. Roden DM, Harrington RA, Poppas A, et al. Considerations for drug interactions on QTc in exploratory COVID-19 (Coronavirus Disease 2019) treatment. *Circulation* 2020. [Epub ahead of print] PMID: 32267762

126. Borba MGS, Val FFA, Sampaio VS, et al. Effect of high vs low doses of chloroquine diphosphate as adjunctive therapy for patients hospitalized with Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2) infection: a randomized clinical trial. *JAMA Netw Open* 2020;3:e208857. PMID: 32330277
127. Tang W, Cao Z, Han M, et al. Hydroxychloroquine in patients with COVID-19: an open-label, randomized, controlled trial. Pre-print. Retrieved April 16, 2020. Available on the World Wide Web at: <https://www.medrxiv.org/content/10.1101/2020.04.10.20060558v1>
128. American Society of Hematology (ASH). COVID-19 and pulmonary embolism: frequently asked questions Version 1.0. Retrieved April 20, 2020. Available on the World Wide Web at: <https://www.hematology.org/covid-19/covid-19-and-pulmonary-embolism>
129. Avigan (favipiravir) tablets package insert [English translation]. Tokyo, Japan: Taisho Toyama Pharmaceutical Co., Ltd.; 2017 Nov. Accessed 2020 Apr 14. Available at: https://www.cdc.gov.tw/File/Get/ht8jUjB_MI-aKnlwstzwv
130. Jakafi (ruxolitinib) package insert. Wilmington, DE: Incyte Corporation; 2020 Jan.
131. Xeljanz (tofacitinib) package insert. New York, NY: Pfizer, Inc.; 2019 Dec.
132. ClinicalTrials.gov. National Institutes of Health (NIH) U.S. National Library of Medicine ClinicalTrials.gov website <https://clinicaltrials.gov>.
133. COVID-19 Treatment Guidelines Panel. Coronavirus Diseases 2019 (COVID-19) Treatment Guidelines. National Institutes of Health. Accessed May 12, 2020. Available at on the World Wide Web at: <https://covid19treatmentguidelines.nih.gov>
134. Magagnoli J, Narendran S, Pereira F, et al. Outcomes of hydroxychloroquine usage in United States veterans hospitalized with Covid-19. Pre-print. Retrieved April 22, 2020. Available on the World Wide Web at: <https://www.medrxiv.org/content/10.1101/2020.04.16.20065920v1>
135. Ye M, Fu D, Ren Y, et al. Treatment with convalescent plasma for COVID-19 patients in Wuhan, China. *J Med Virol*. 2020. [Epub ahead of print] PMID: 32293713
136. Wang J, Hajizadeh N, Moore EE, et al. Tissue plasminogen activator (tPA) treatment for COVID-19 associated acute respiratory distress syndrome (ARDS): a case series. *J Thromb Haemost*. 2020. [Epub ahead of print] PMID: 32267998
137. Activase (alteplase) package insert. South San Francisco, CA: Genentech, Inc.; 2018 Feb.
138. Beth Israel Deaconess Medical Center. BIDMC launches clinical trial to assess common anti-clotting medication for treatment of COVID-19-related respiratory failure. Retrieved April 22, 2020. Available on the World Wide Web at <https://www.bidmc.org/about-bidmc/news/2020/04/covid-19-anti-clotting-medication>
139. IRCCS San Raffaele. Defibrotide in COVID-19 pneumonia (DEFI-VID19). Retrieved April 17, 2020. Available on the World Wide Web at: <https://clinicaltrials.gov/ct2/show/NCT04335201?term=defibrotide&cond=COVID&draw=2&rank=2>
140. Fundacion para la Formacion e Investigacion Sanitarias de la Region de Murcia. Defibrotide as prevention and treatment of respiratory distress and cytokine release syndrome of COVID 19 (DEFACOVID). Retrieved April 17, 2020. Available on the World Wide Web at: <https://clinicaltrials.gov/ct2/show/NCT04348383?term=defibrotide&cond=COVID&draw=2&rank=1>
141. Food and Drug Administration. FDA Drug Safety Communication: Hydroxychloroquine or chloroquine for COVID-19: FDA cautions against use outside of the hospital setting or a clinical trial due to risk of heart rhythm problems. Retrieved April 23, 2020. Available on the World Wide Web at: https://www.fda.gov/safety/medical-product-safety-information/hydroxychloroquine-or-chloroquine-covid-19-drug-safety-communication-fda-cautions-against-use?utm_campaign=FDA%20MedWatch%20

142. Cai Q, Yang M, Liu D, et al. Experimental treatment with favipiravir for COVID-19: An open-label control study. *Engineering*. <https://doi.org/10.1016/j.eng.2020.03.007>
143. Chen C, Zhang Y, Huang J, et al. Favipiravir versus Arbidol for COVID-19: A Randomized Clinical Trial. Pre-print. Retrieved April 27, 2020. Available on the World Wide Web at: <https://doi.org/10.1101/2020.03.17.20037432>
144. Cantini F, Niccoli L, Matarrese D, et al. Baricitinib therapy in COVID-19: A pilot study on safety and clinical impact. *J Infect* 2020 April 22;S0163-4453(20)30228-0. [Epub Ahead of Print] PMID: 32333918
145. NIAID Office of Communications. NIH clinical trial shows remdesivir accelerates recovery from advanced COVID-19. National Institutes of Health Web site. Accessed April 29, 2020. Available at: <https://www.niaid.nih.gov/news-events/nih-clinical-trial-shows-remdesivir-accelerates-recovery-advanced-covid-19>
146. Food and Drug Administration (FDA). Fact sheet for health care providers emergency use authorization (EUA) of remdesivir (GS-5734). Retrieved May 4, 2020. Available on the World Wide Web at: <https://www.fda.gov/media/137566/download>
147. Ilaris (canakinumab) package insert. East Hanover, NJ: Novartis Pharmaceuticals Corporation; 2016 Dec.
148. Cavalli G, De Luca G, Campochiaro C, et al. Interleukin-1 blockade with high-dose anakinra in patients with COVID-19, acute respiratory distress syndrome, and hyperinflammation: a retrospective cohort study. *Lancet Rheumatol* 2020; [https://doi.org/10.1016/S2665-9913\(20\)30127-2](https://doi.org/10.1016/S2665-9913(20)30127-2) [Epub ahead of print]
149. Geleris J, Sun Y, Platt J, et al. Observational study of hydroxychloroquine in hospitalized patients with COVID-19. *N Engl J Med* 2020 May 7. [Epub ahead of print] PMID: 32379955
150. Wang Y, Zhang D, Du G, et al. Remdesivir in adults with severe COVID-19: a randomized, double-blind, placebo-controlled, multicenter trial. *The Lancet*. 2020. Available at: [https://www.thelancet.com/journals/lancet/article/PIIS0140-6736\(20\)31022-9/fulltext#seccestitle10](https://www.thelancet.com/journals/lancet/article/PIIS0140-6736(20)31022-9/fulltext#seccestitle10) [Epub ahead of print]
151. Rosenberg ES, Dufort EM, Udo T, et al. Association of treatment with hydroxychloroquine or azithromycin with in-hospital mortality in patients with COVID-19 in New York state. *JAMA* 2020 [Epub ahead of print] PMID: 32392282